

Cytokeratin 20, Biotinylated
Peptide-affinity purified goat antibody
Catalog # AF3856b

Specification

Cytokeratin 20, Biotinylated - Product Information

Application	WB
Primary Accession	P35900
Other Accession	NP_061883.1 , 54474
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	48487

Cytokeratin 20, Biotinylated - Additional Information

Gene ID 54474

Other Names

Keratin, type I cytoskeletal 20,
Cytokeratin-20, CK-20, Keratin-20, K20,
Protein IT, KRT20

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

Cytokeratin 20, Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

Cytokeratin 20, Biotinylated - Protein Information

Name KRT20

Function



Biotinylated EB11745 (1µg/ml) staining of Human Colon lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker

Cytokeratin 20, Biotinylated - References

SATB2 in combination with cytokeratin 20 identifies over 95% of all colorectal carcinomas. Magnusson K, de Wit M, Brennan DJ, Johnson LB, McGee SF, Lundberg E, Naicker K, Klinger R, Kampf C, Asplund A, Wester K, Gry M, Bjartell A, Gallagher WM, Rexhepaj E, Kilpinen S, Kallioniemi OP, Belt E, Goos J, Meijer G, Birgisson H, Glimelius B, Borrebaeck AM J Surg Pathol. 2011 Jul;35(7):937-48. PMID: 21677534

Plays a significant role in maintaining keratin filament organization in intestinal epithelia. When phosphorylated, plays a role in the secretion of mucin in the small intestine (By similarity).

Cellular Location

Cytoplasm

Tissue Location

Expressed predominantly in the intestinal epithelium. Expressed in luminal cells of colonic mucosa. Also expressed in the Merkel cells of keratinized oral mucosa; specifically at the tips of some rete ridges of the gingival mucosa, in the basal layer of the palatal mucosa and in the taste buds of lingual mucosa

Cytokeratin 20, Biotinylated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)