

Proximal Nephrogenic Antigen (Renal Cell Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone PN-15]
 Catalog # AH12554

Specification

Proximal Nephrogenic Antigen (Renal Cell Marker) Antibody - With BSA and Azide - Product Information

Application	,1,2,3,4,
Primary Accession	Q16790
Other Accession	768 , 63287
Reactivity	Human, Horse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2b, kappa
Calculated MW	55kDa KDa

Proximal Nephrogenic Antigen (Renal Cell Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 768

Other Names

Carbonic anhydrase 9, 4.2.1.1, Carbonate dehydratase IX, Carbonic anhydrase IX, CA-IX, CAIX, Membrane antigen MN, P54/58N, Renal cell carcinoma-associated antigen G250, RCC-associated antigen G250, pMW1, CA9, G250, MN

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions


Proximal Nephrogenic Antigen (Renal Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Proximal Nephrogenic Antigen (Renal Cell Marker) Antibody - With BSA and Azide - Protein Information

Name CA9

Synonyms G250, MN

Function



Proximal Nephrogenic Antigen (Renal Cell Marker) Antibody - With BSA and Azide - Background

Recognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowman's capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This MAb may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.

Proximal Nephrogenic Antigen (Renal Cell Marker) Antibody - With BSA and Azide - References

Reversible hydration of carbon dioxide. Participates in pH regulation. May be involved in the control of cell proliferation and transformation. Appears to be a novel specific biomarker for a cervical neoplasia.

Yoshida SO, et. al. Cancer Research, 1989, 49(7):1802-9

Cellular Location

Nucleus. Nucleus, nucleolus. Cell membrane; Single-pass type I membrane protein. Cell projection, microvillus membrane; Single-pass type I membrane protein. Note=Found on the surface microvilli and in the nucleus, particularly in nucleolus

Tissue Location

Expressed primarily in carcinoma cells lines. Expression is restricted to very few normal tissues and the most abundant expression is found in the epithelial cells of gastric mucosa

Proximal Nephrogenic Antigen (Renal Cell Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)