

SUZ12 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI10063

Specification

SUZ12 antibody - middle region - Product Information

Application	CHIP, IHC, WB
Primary Accession	Q15022
Other Accession	Q15022 , NP_056170 , NM_015355
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rabbit, Chicken, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83 kDa KDa

SUZ12 antibody - middle region - Additional Information

Gene ID 23512


Alias Symbol **CHET9, JJAZ1, KIAA0160**

Other Names


Polycomb protein SUZ12, Chromatin precipitated E2F target 9 protein, ChET 9 protein, Joined to JAZF1 protein, Suppressor of zeste 12 protein homolog, SUZ12, CHET9, JJAZ1, KIAA0160

Target/Specificity

A chromosomal aberration involving SUZ12 may be a cause of endometrial stromal tumors. Translocation t (7, 17)(p15, q21) with JAZF1 generates the JAZF1-SUZ12 oncogene consisting of the N-terminus part of JAZF1 and the C-terminus part of SUZ12. It is frequently found in all cases of endometrial stromal tumors, except in endometrial stromal sarcomas, where it is rarer. This zinc finger gene has been identified at the breakpoints of a recurrent chromosomal translocation reported in



SUZ12 antibody - middle region (AI10063) in HCT116 cells using Chromatin Immunoprecipitation (ChIP) Using SUZ12 antibody - middle region (AI10063) and HCT116 Cells



SUZ12 antibody - middle region (AI10063) in Human Placenta cells using Immunohistochemistry Placenta

endometrial stromal sarcoma.

Recombination of these breakpoints results in the fusion of this gene and JAZF1. The protein encoded by this gene contains a zinc finger domain in the C terminus of the coding region. The specific function of this gene has not yet been determined.

Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SUZ12 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

SUZ12 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.


SUZ12 antibody - middle region - Protein Information

Name SUZ12

Synonyms CHET9, JJAZ1, KIAA0160

Function

Polycomb group (PcG) protein. Component of the PRC2 complex, which methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to transcriptional repression of the affected target gene (PubMed:15225548, PubMed:15231737, PubMed:15385962, PubMed:16618801, PubMed:<a href="http://www.uniprot.org/ci



SUZ12 antibody - middle region (AI10063) in Human HeLa cells using Western Blot
WB Suggested Anti-SUZ12 Antibody Titration: 0.2-1 µg/ml
Positive Control: Hela cell lysate
SUZ12 is strongly supported by BioGPS gene expression data to be expressed in Human HeLa cells

SUZ12 antibody - middle region - Background

This is a rabbit polyclonal antibody against SUZ12. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).

tations/17344414"
target="_blank">17344414,
PubMed:<a href="http://www.uniprot.org/ci
tations/18285464"
target="_blank">18285464,
PubMed:<a href="http://www.uniprot.org/ci
tations/28229514"
target="_blank">28229514,
PubMed:<a href="http://www.uniprot.org/ci
tations/29499137"
target="_blank">29499137,
PubMed:<a href="http://www.uniprot.org/ci
tations/31959557"
target="_blank">31959557). The
PRC2 complex may also serve as a
recruiting platform for DNA
methyltransferases, thereby linking two
epigenetic repression systems (PubMed:<a
href="http://www.uniprot.org/citations/1243
5631" target="_blank">12435631,
PubMed:<a href="http://www.uniprot.org/ci
tations/12351676"
target="_blank">12351676,
PubMed:<a href="http://www.uniprot.org/ci
tations/15385962"
target="_blank">15385962,
PubMed:<a href="http://www.uniprot.org/ci
tations/15099518"
target="_blank">15099518,
PubMed:<a href="http://www.uniprot.org/ci
tations/15225548"
target="_blank">15225548,
PubMed:<a href="http://www.uniprot.org/ci
tations/15684044"
target="_blank">15684044,
PubMed:<a href="http://www.uniprot.org/ci
tations/16431907"
target="_blank">16431907,
PubMed:<a href="http://www.uniprot.org/ci
tations/18086877"
target="_blank">18086877,
PubMed:<a href="http://www.uniprot.org/ci
tations/18285464"
target="_blank">18285464). Genes
repressed by the PRC2 complex include
HOXC8, HOXA9, MYT1 and CDKN2A
(PubMed:<a href="http://www.uniprot.org/c
itations/15231737"
target="_blank">15231737,
PubMed:<a href="http://www.uniprot.org/ci
tations/16618801"
target="_blank">16618801,
PubMed:<a href="http://www.uniprot.org/ci
tations/17200670"
target="_blank">17200670,
PubMed:<a href="http://www.uniprot.org/ci
tations/31959557"

target="_blank">31959557).

Cellular Location

Nucleus Note=Localizes to chromatin as part of the PRC2 complex

Tissue Location

Overexpressed in breast and colon cancer.

**SUZ12 antibody - middle region -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)