

## TAF9 antibody - N-terminal region

### Rabbit Polyclonal Antibody

Catalog # AI10686

### Specification

#### TAF9 antibody - N-terminal region - Product Information

Application **CHIP, IHC, WB**  
Primary Accession [Q9Y3D8](#)  
Other Accession [NM\\_001015891](#),  
[NP\\_001015891](#)  
Reactivity **Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine**  
Host **Rabbit**  
Clonality **Polyclonal**  
Calculated MW **19kDa KDa**

#### TAF9 antibody - N-terminal region - Additional Information

**Gene ID** 102157402


**Alias Symbol** **AK6, CIP, CINAP, TAF2G, AD-004, hCINAP, CGI-137, TAFII31, TAFII32, MGC:1603, MGC:3647, MGC:5067, TAFIID32**

#### Other Names


Adenylate kinase isoenzyme 6  
{ECO:0000255|HAMAP-Rule:MF\_03173},  
AK6  
{ECO:0000255|HAMAP-Rule:MF\_03173},  
2.7.4.3  
{ECO:0000255|HAMAP-Rule:MF\_03173},  
Adrenal gland protein AD-004,  
Coilin-interacting nuclear ATPase protein  
{ECO:0000255|HAMAP-Rule:MF\_03173},  
hCINAP, Dual activity adenylate  
kinase/ATPase  
{ECO:0000255|HAMAP-Rule:MF\_03173},  
AK/ATPase  
{ECO:0000255|HAMAP-Rule:MF\_03173},  
AK6  
{ECO:0000255|HAMAP-Rule:MF\_03173}

#### Format


Liquid. Purified antibody supplied in 1x PBS



Quiescent human colon carcinoma HCT116 cultures were treated with 10% FBS for three time points (0, 15, 30min) or (0, 30, 60min) were used in Matrix-ChIP and real-time PCR assays at EGR1 gene (Exon1) and 15kb upstream site.



Human Skin



WB Suggested Anti-TAF9 Antibody Titration:  
0.125µg/ml  
ELISA Titer: 1:312500

buffer with 0.09% (w/v) sodium azide and 2% sucrose.

#### **Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-TAF9 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

#### **Precautions**

TAF9 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

#### **TAF9 antibody - N-terminal region - Protein Information**

##### **Name** AK6

{ECO:0000255|HAMAP-Rule:MF\_03173}

##### **Function**

Broad-specificity nucleoside monophosphate (NMP) kinase that catalyzes the reversible transfer of the terminal phosphate group between nucleoside triphosphates and monophosphates. AMP and dAMP are the preferred substrates, but CMP and dCMP are also good substrates. IMP is phosphorylated to a much lesser extent. All nucleoside triphosphates ATP, GTP, UTP, CTP, dATP, dCTP, dGTP, and TTP are accepted as phosphate donors. CTP is the best phosphate donor, followed by UTP, ATP, GTP and dCTP. May have a role in nuclear energy homeostasis. Has also ATPase activity. May be involved in regulation of Cajal body (CB) formation.

##### **Cellular Location**

Nucleus, nucleoplasm. Nucleus, Cajal body. Note=Displays widespread diffuse nucleoplasmic distribution but not detected in nucleoli. Detected in Cajal bodies but not in all cells

##### **Tissue Location**

Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, chorionic villi and the central nervous system.

#### **Positive Control: Human Small Intestine**

#### **TAF9 antibody - N-terminal region - References**

Evans,S.C., et al., (1999) 57 (1), 182-183  
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

#### **TAF9 antibody - N-terminal region -**

## Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)