

Thrb antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11090

Specification

Thrb antibody - N-terminal region - Product Information

Application **CHIP, WB**
Primary Accession [P37244](#)
Other Accession [NM_009380](#),
[NP_033406](#)
Reactivity **Human, Mouse, Rat, Pig, Horse, Bovine**
Predicted **Mouse, Rat**
Host **Rabbit**
Clonality **Polyclonal**
Calculated MW **52kDa KDa**

Thrb antibody - N-terminal region - Additional Information

Alias Symbol **Nr1a2, T3R[b], T3Rbeta, Thrb1, Thrb2, c-erbAbeta**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage


Add 100 ul of distilled water. Final anti-Thrb antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Thrb antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Thrb antibody - N-terminal region - Protein Information

Thrb antibody - N-terminal region - Protocols



See Other Applications for more information.

Application: ChIP
Sample Type: mouse liver tissue Chromatin
Used: 100ug tissue Antibody Used: 10ug
Image
Submitted by: Joanna DiSpirito University of Pennsylvania




See Other Applications for more information.

Application: ChIP
Sample Type: mouse liver tissue Chromatin
Used: 100ug tissue Antibody Used: 10ug
Image
Submitted by: Joanna DiSpirito University of Pennsylvania

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



WB Suggested Anti-Thrb Antibody Titration:
1.25 μ g/ml
Positive Control: SP2/0 cell lysate

Thrb antibody - N-terminal region - References

Furuya, F., (2006) Proc. Natl. Acad. Sci. U.S.A. 103 (6), 1780-1785
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.