

SMARCA1 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11216

Specification

SMARCA1 antibody - N-terminal region - Product Information

Application	CHIP, WB
Primary Accession	Q8BS67
Other Accession	NM_053123 , BAC28931
Reactivity	Mouse, Rat
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa

SMARCA1 antibody - N-terminal region - Additional Information

Alias Symbol **Snf2l,
5730494M04Rik**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-SMARCA1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

SMARCA1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


SMARCA1 antibody - N-terminal region - Protein Information

SMARCA1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.



Chromatin Immunoprecipitation (ChIP) Using SMARCA1 antibody - N-terminal region (AI11216) and HCT116 Cells



WB Suggested Anti-SMARCA1 Antibody Titration: 1.25µg/ml
ELISA Titer: 1:312500
Positive Control: NIH/3T3 cell lysate

SMARCA1 antibody - N-terminal region - References

Okazaki,Y., et al., (2002) Nature 420 (6915), 563-573
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)