

IL11RA antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12686

Specification

IL11RA antibody - middle region - Product Information

Application	WB
Primary Accession	Q14626
Other Accession	NM_004512 , NP_004503
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rabbit, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa

IL11RA antibody - middle region - Additional Information

Gene ID 3590

Alias Symbol **MGC2146, CRSDA**
Other Names
Interleukin-11 receptor subunit alpha, IL-11
receptor subunit alpha, IL-11R subunit
alpha, IL-11R-alpha, IL-11RA, IL11RA

Format


Liquid. Purified antibody supplied in 1x PBS
buffer with 0.09% (w/v) sodium azide and
2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final
anti-IL11RA antibody concentration is 1
mg/ml in PBS buffer with 2% sucrose. For
longer periods of storage, store at 20°C.
Avoid repeat freeze-thaw cycles.

Precautions

IL11RA antibody - middle region is for
research use only and not for use in
diagnostic or therapeutic procedures.



WB Suggested Anti-IL11RA Antibody
Titration: 0.2-1 µg/ml
ELISA Titer: 1:12500
Positive Control: Human Muscle

IL11RA antibody - middle region - References

Nakayama,T.,(2007)Int.J.Oncol.30(4),825-833R
econstitutionandStorage:Forshorttermuse,store
at-8Cupto1week.Forlongtermstorage,storeat-
20Cinsmallaliquotstopreventfreeze-thawcycles.

IL11RA antibody - middle region - Protein Information

Name IL11RA ([HGNC:5967](#))

Function

Receptor for interleukin-11 (IL11). The receptor systems for IL6, LIF, OSM, CNTF, IL11 and CT1 can utilize IL6ST for initiating signal transmission. The IL11/IL11RA/IL6ST complex may be involved in the control of proliferation and/or differentiation of skeletogenic progenitor or other mesenchymal cells (Probable). Essential for the normal development of craniofacial bones and teeth. Restricts suture fusion and tooth number.

Cellular Location

[Interleukin-11 receptor subunit alpha]:
Membrane; Single-pass type I membrane protein [Isoform HCR2]: Secreted

Tissue Location

Expressed in a number of cell lines, including the myelogenous leukemia cell line K-562, the megakaryocytic leukemia cell line M-07e, the erythroleukemia cell line TF-1, and the osteosarcoma cell lines, MG-63 and SaOS-2 (PubMed:7670098). Also expressed in normal and malignant prostate epithelial cell lines. Expression levels are increased in prostate carcinoma.

IL11RA antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)