


HHIPL1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14475

Specification

HHIPL1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q96JK4
Other Accession	NM_001127258 , NP_001120730
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Rat, Rabbit, Pig, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85kDa KDa



WB Suggested Anti-HHIPL1 Antibody
Titration: 0.2-1 µg/ml
ELISA Titer: 1:62500
Positive Control: MDA-MB-435S cell lysate

HHIPL1 antibody - C-terminal region - Additional Information

Gene ID 84439

Alias Symbol **KIAA1822,**
UNQ9245

Other Names

HHIP-like protein 1, HHIPL1, HHIP2,
KIAA1822

Format

Liquid. Purified antibody supplied in 1x PBS
buffer with 0.09% (w/v) sodium azide and
2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final
anti-HHIPL1 antibody concentration is 1
mg/ml in PBS buffer with 2% sucrose. For
longer periods of storage, store at 20°C.
Avoid repeat freeze-thaw cycles.

Precautions

HHIPL1 antibody - C-terminal region is for
research use only and not for use in
diagnostic or therapeutic procedures.

HHIPL1 antibody - C-terminal region - References

Clark H.F., et al. Genome Res.
13:2265-2270(2003).
Heilig R., et al. Nature 421:601-607(2003).
Nagase T., et al. DNA Res. 8:85-95(2001).
Katoh Y., et al. Int. J. Mol. Med.
17:391-395(2006).

HHIPL1 antibody - C-terminal region - Protein Information

Name HHIPL1

Synonyms HHIP2, KIAA1822

Cellular Location
Secreted.

HHIPL1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)