

FMR1 / FMRP Antibody (C-Terminus)
Goat Polyclonal Antibody
Catalog # ALS13094

Specification

FMR1 / FMRP Antibody (C-Terminus) - Product Information

Application	WB, IHC
Primary Accession	Q06787
Reactivity	Human, Mouse, Rat, Rabbit, Hamster, Monkey, Pig
Host	Goat
Clonality	Polyclonal
Calculated MW	71kDa KDa

FMR1 / FMRP Antibody (C-Terminus) - Additional Information

Gene ID 2332

Other Names

Fragile X mental retardation protein 1, FMRP, Protein FMR-1, FMR1

Target/Specificity

Human FMR1.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

FMR1 / FMRP Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

FMR1 / FMRP Antibody (C-Terminus) - Protein Information


Name FMR1 ([HGNC:3775](#))

Function

Multifunctional polyribosome-associated RNA-binding protein that plays a central role in neuronal development and synaptic plasticity through the regulation of alternative mRNA splicing, mRNA stability, mRNA dendritic transport and postsynaptic



Antibody (0.1 ug/ml) staining of KELLY cell lysate (35 ug protein in RIPA buffer).



Anti-FMR1 antibody IHC of human placenta.

FMR1 / FMRP Antibody (C-Terminus) - Background

Translation repressor. Component of the CYFIP1-EIF4E- FMR1 complex which binds to the mRNA cap and mediates translational repression. In the CYFIP1-EIF4E-FMR1 complex this subunit mediates translation repression (By similarity). RNA- binding protein that plays a role in intracellular RNA transport and in the regulation of translation of target mRNAs. Associated with polysomes. May play a role in the transport of mRNA from the nucleus to the

local protein synthesis of a subset of mRNAs (PubMed:16631377, PubMed:18653529, PubMed:19166269, PubMed:23235829, PubMed:25464849). Plays a role in the alternative splicing of its own mRNA (PubMed:18653529). Plays a role in mRNA nuclear export (By similarity). Together with export factor NXF2, is involved in the regulation of the NXF1 mRNA stability in neurons (By similarity). Stabilizes the scaffolding postsynaptic density protein DLG4/PSD-95 and the myelin basic protein (MBP) mRNAs in hippocampal neurons and glial cells, respectively; this stabilization is further increased in response to metabotropic glutamate receptor (mGluR) stimulation (By similarity). Plays a role in selective delivery of a subset of dendritic mRNAs to synaptic sites in response to mGluR activation in a kinesin-dependent manner (By similarity). Plays a role as a repressor of mRNA translation during the transport of dendritic mRNAs to postsynaptic dendritic spines (PubMed:11532944, PubMed:11157796, PubMed:12594214, PubMed:23235829). Component of the CYFIP1-EIF4E-FMR1 complex which blocks cap-dependent mRNA translation initiation (By similarity). Represses mRNA translation by stalling ribosomal translocation during elongation (By similarity). Reports are contradictory with regards to its ability to mediate

cytoplasm. Binds strongly to poly(G), binds moderately to poly(U) but shows very little binding to poly(A) or poly(C).

FMR1 / FMRP Antibody (C-Terminus) - References

Verkerk A.J.M.H.,et al.Cell 65:905-914(1991).
Verkerk A.J.M.H.,et al.Hum. Mol. Genet. 2:399-404(1993).
Verkerk A.J.H.M.,et al.Hum. Mol. Genet. 2:1348-1348(1993).
Corominas R.,et al.Nat. Commun. 5:3650-3650(2014).
Ross M.T.,et al.Nature 434:325-337(2005).

translation inhibition of MBP mRNA in oligodendrocytes (PubMed:23891804). Also involved in the recruitment of the RNA helicase MOV10 to a subset of mRNAs and hence regulates microRNA (miRNA)-mediated translational repression by AGO2 (PubMed:14703574, PubMed:17057366, PubMed:25464849). Facilitates the assembly of miRNAs on specific target mRNAs (PubMed:17057366). Plays also a role as an activator of mRNA translation of a subset of dendritic mRNAs at synapses (PubMed:19097999, PubMed:19166269). In response to mGluR stimulation, FMR1-target mRNAs are rapidly derepressed, allowing for local translation at synapses (By similarity). Binds to a large subset of dendritic mRNAs that encode a myriad of proteins involved in pre- and postsynaptic functions (PubMed:7692601, PubMed:11719189, PubMed:11157796, PubMed:12594214, PubMed:17417632, PubMed:23235829, PubMed:24448548). Binds to 5'-ACU[GU]-3' and/or 5'-[AU]GGA-3' RNA

consensus sequences within mRNA targets, mainly at coding sequence (CDS) and 3'-untranslated region (UTR) and less frequently at 5'-UTR (PubMed:23235829). Binds to intramolecular G-quadruplex structures in the 5'- or 3'-UTRs of mRNA targets (PubMed:11719189, PubMed:18579868, PubMed:25464849, PubMed:25692235). Binds to G-quadruplex structures in the 3'-UTR of its own mRNA (PubMed:7692601, PubMed:11532944, PubMed:12594214, PubMed:15282548, PubMed:18653529). Binds also to RNA ligands harboring a kissing complex (kc) structure; this binding may mediate the association of FMR1 with polyribosomes (PubMed:15805463). Binds mRNAs containing U-rich target sequences (PubMed:12927206). Binds to a triple stem-loop RNA structure, called Sod1 stem loop interacting with FMRP (SoSLIP), in the 5'-UTR region of superoxide dismutase SOD1 mRNA (PubMed:19166269). Binds to the dendritic, small non-coding brain cytoplasmic RNA 1 (BC1); which may increase the association of the CYFIP1-EIF4E-FMR1 complex to FMR1 target mRNAs at synapses (By similarity).

Associates with export factor NXF1 mRNA-containing ribonucleoprotein particles (mRNPs) in a NXF2-dependent manner (By similarity). Binds to a subset of miRNAs in the brain (PubMed:14703574, PubMed:17057366). May associate with nascent transcripts in a nuclear protein NXF1-dependent manner (PubMed:18936162). In vitro, binds to RNA homomer; preferentially on poly(G) and to a lesser extent on poly(U), but not on poly(A) or poly(C) (PubMed:7688265, PubMed:7781595, PubMed:12950170, PubMed:15381419, PubMed:8156595). Moreover, plays a role in the modulation of the sodium-activated potassium channel KCNT1 gating activity (PubMed:20512134). Negatively regulates the voltage-dependent calcium channel current density in soma and presynaptic terminals of dorsal root ganglion (DRG) neurons, and hence regulates synaptic vesicle exocytosis (By similarity). Modulates the voltage-dependent calcium channel CACNA1B expression at the plasma membrane by targeting the channels for proteosomal degradation (By similarity). Plays a role in regulation of MAP1B-dependent microtubule dynamics during neuronal development (By similarity). Recently, has been shown to play a translation-independent role in the modulation of presynaptic action potential (AP) duration and neurotransmitter release via large- conductance calcium-activated potassium (BK) channels in hippocampal and cortical excitatory neurons (PubMed:<a href="http://www.uniprot.org/citations/2556

1520" target="_blank">25561520).
Finally, FMR1 may be involved in the control of DNA damage response (DDR) mechanisms through the regulation of ATR-dependent signaling pathways such as histone H2AX/H2A.x and BRCA1 phosphorylations (PubMed:24813610).

Cellular Location

Nucleus, Nucleus, nucleolus. Chromosome, centromere

{ECO:0000250|UniProtKB:P35922}.

Chromosome

{ECO:0000250|UniProtKB:P35922}.

Cytoplasm. Cytoplasm, perinuclear region.

Cytoplasm, Cytoplasmic ribonucleoprotein granule. Perikaryon. Cell projection, neuron projection. Cell projection, axon

{ECO:0000250|UniProtKB:P35922}. Cell

projection, dendrite

{ECO:0000250|UniProtKB:P35922}. Cell

projection, dendritic spine

{ECO:0000250|UniProtKB:P35922}. Cell

junction, synapse, synaptosome

{ECO:0000250|UniProtKB:P35922}. Cell

projection, growth cone. Cell projection,

filopodium tip

{ECO:0000250|UniProtKB:P35922}. Cell

junction, synapse

{ECO:0000250|UniProtKB:P35922}. Cell

junction, synapse, postsynaptic cell

membrane

{ECO:0000250|UniProtKB:P35922}. Cell

junction, synapse, presynaptic cell

membrane

{ECO:0000250|UniProtKB:P35922}. Cell

membrane

{ECO:0000250|UniProtKB:P35922}.

Cytoplasm, Stress granule.

Note=Colocalizes with H2AX/H2A.x in

pericentromeric heterochromatin in

response to DNA damaging agents (By

similarity). Localizes on meiotic

pachytene-stage chromosomes (By

similarity). Forms nuclear foci representing

sites of ongoing DNA replication in response

to DNA damaging agents (By similarity)

Shuttles between nucleus and cytoplasm in

a XPO1/CRM1-dependent manner

(PubMed:10196376). Localizes to

cytoplasmic ribonucleoprotein granules,

also referred to as messenger

ribonucleoprotein particles or mRNPs, along

dendrites and dendritic spines

(PubMed:9659908, PubMed:14532325)

FMR1-containing cytoplasmic granules colocalize to F-actin-rich structures, including filopodium, spines and growth cone during the development of hippocampal neurons (By similarity). FMR1-containing cytoplasmic granules are transported out of the soma along axon and dendrite to synaptic contacts in a microtubule- and kinesin-dependent manner (PubMed:12417734, PubMed:15380484). Colocalizes with CACNA1B in the cytoplasm and at the cell membrane of neurons (By similarity). Colocalizes with CYFIP1, CYFIP2, NXF2 and ribosomes in the perinuclear region (By similarity). Colocalizes with CYFIP1 and EIF4E in dendrites and probably at synapses (By similarity). Colocalizes with FXR1, kinesin, 60S acidic ribosomal protein RPLP0 and SMN in cytoplasmic granules in the soma and neurite cell processes (PubMed:12417734, PubMed:18093976, PubMed:16636078). Colocalizes with FXR1 and FXR2 in discrete granules, called fragile X granules (FXGs), along axon and presynaptic compartments (By similarity). Colocalizes with TDRD3 in cytoplasmic stress granules (SGs) in response to various cellular stress (PubMed:18632687, PubMed:18664458, PubMed:16636078) {ECO:0000250|UniProtKB:P35922, ECO:0000250|UniProtKB:Q80WE1, ECO:0000269|PubMed:10196376, ECO:0000269|PubMed:12417734, ECO:0000269|PubMed:14532325, ECO:0000269|PubMed:15380484, ECO:0000269|PubMed:16636078, ECO:0000269|PubMed:18093976, ECO:0000269|PubMed:18632687, ECO:0000269|PubMed:18664458, ECO:0000269|PubMed:9659908} [Isoform 9]: Cytoplasm [Isoform 11]: Nucleus. Nucleus, Cajal body

Tissue Location

Expressed in the brain, cerebellum and testis (PubMed:8401578). Also expressed in epithelial tissues (PubMed:8401578). Expressed in mature oligodendrocytes (OLGs) (PubMed:23891804). Expressed in fibroblast (PubMed:24204304). Expressed in neurons, Purkinje cells and spermatogonias (at protein level) (PubMed:8401578). Expressed in brain, testis and placenta (PubMed:8504300). Expressed in neurons and lymphocytes (PubMed:8504300).

**FMR1 / FMRP Antibody (C-Terminus) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)