

NR2F1 / Coup-TF Antibody (aa1-44)
Rabbit Polyclonal Antibody
Catalog # ALS16406

Specification

NR2F1 / Coup-TF Antibody (aa1-44) - Product Information

Application	WB, ICC, IHC
Primary Accession	P10589
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46kDa KDa

NR2F1 / Coup-TF Antibody (aa1-44) - Additional Information

Gene ID 7025

Other Names

COUP transcription factor 1, COUP-TF1, COUP transcription factor I, COUP-TF I, Nuclear receptor subfamily 2 group F member 1, V-erbA-related protein 3, EAR-3, NR2F1, EAR3, ERBAL3, TFCOUP1

Target/Specificity

Human NR2F1 / Coup-TF

Reconstitution & Storage

Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

Precautions

NR2F1 / Coup-TF Antibody (aa1-44) is for research use only and not for use in diagnostic or therapeutic procedures.


NR2F1 / Coup-TF Antibody (aa1-44) - Protein Information

Name NR2F1

Synonyms EAR3, ERBAL3, TFCOUP1

Function


Coup (chicken ovalbumin upstream promoter) transcription factor binds to the



Western blot of NR2F1 expression in transfected 293T cell line by NR2F1 polyclonal antibody.



Immunofluorescence of paraformaldehyde-fixed HeLa using COUP TF1 antibody at 1:200 dilution.



Human Liver: Formalin-Fixed, Paraffin-Embedded (FFPE)

ovalbumin promoter and, in conjunction with another protein (S300-II) stimulates initiation of transcription. Binds to both direct repeats and palindromes of the 5'-AGGTCA-3' motif. Represses transcriptional activity of LHCG.


Cellular Location

Nucleus.

NR2F1 / Coup-TF Antibody (aa1-44) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Human Skin: Formalin-Fixed,
Paraffin-Embedded (FFPE)

NR2F1 / Coup-TF Antibody (aa1-44) - Background

Coup (chicken ovalbumin upstream promoter) transcription factor binds to the ovalbumin promoter and, in conjunction with another protein (S300-II) stimulates initiation of transcription. Binds to both direct repeats and palindromes of the 5'-AGGTCA-3' motif. Represses transcriptional activity of LHCG.

NR2F1 / Coup-TF Antibody (aa1-44) - References

Wang L.-H., et al. Nature 340:163-166(1989).
Miyajima N., et al. Nucleic Acids Res. 16:11057-11074(1988).
Dressel U., et al. Mol. Cell. Biol. 19:3383-3394(1999).
Zhang Y., et al. J. Biol. Chem. 275:2763-2770(2000).
Zhang Y., et al. Mol. Endocrinol. 15:1891-1905(2001).