

ATG16L1 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM1817b

Specification

ATG16L1 Antibody - Product Information

Application	WB, IHC-P,E
Primary Accession	Q676U5
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1, κ

ATG16L1 Antibody - Additional Information

Gene ID 55054

Other Names

Autophagy-related protein 16-1, APG16-like 1, ATG16L1, APG16L

Target/Specificity

Purified His-tagged ATG16L1 protein(Fragment) was used to produced this monoclonal antibody.

Dilution

WB~~1:60
IHC-P~~1:10~50

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

ATG16L1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

ATG16L1 Antibody - Protein Information



Western blot analysis of anti-ATG16L1 Monoclonal Antibody (Cat.#AM1817b) by ATG16L1 recombinant protein (Fragment). ATG16L1 (Fragment) protein (arrow) was detected using the purified Mab. (1:2000)



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with ATG16L1 Monoclonal Antibody (Cat.#AM1817b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

ATG16L1 Antibody - Background

Name ATG16L1

Synonyms APG16L

Function

Plays an essential role in autophagy: interacts with ATG12- ATG5 to mediate the conjugation of phosphatidylethanolamine (PE) to LC3 (MAP1LC3A, MAP1LC3B or MAP1LC3C), to produce a membrane-bound activated form of LC3 named LC3-II. Thereby, controls the elongation of the nascent autophagosomal membrane

(PubMed:24553140,
PubMed:23376921,
PubMed:24954904,
PubMed:27273576,
PubMed:23392225).

Regulates mitochondrial antiviral signaling (MAVS)-dependent type I interferon (IFN-I) production (PubMed:25645662).

Negatively regulates NOD1- and NOD2-driven inflammatory cytokine response (PubMed:24238340). Instead, promotes with NOD2 an autophagy-dependent antibacterial pathway (PubMed:20637199).

Plays a role in regulating morphology and function of Paneth cell (PubMed:18849966).

Cellular Location

Cytoplasm. Preautophagosomal structure membrane; Peripheral membrane protein.

Note=Recruited to omegasomes membranes by WIPI2 Omegasomes are endoplasmic reticulum connected structures at the origin of preautophagosomal structures. Localized to preautophagosomal structure (PAS) where it is involved in the

The protein encoded by this gene is part of a large protein complex that is necessary for autophagy, the major process by which intracellular components are targeted to lysosomes for degradation. Defects in this gene are a cause of susceptibility to inflammatory bowel disease type 10 (IBD10). Several transcript variants encoding different isoforms have been found for this gene.

ATG16L1 Antibody - References

Age at onset in Huntington's disease is modified by the autophagy pathway: implication of the V471A polymorphism in Atg7. Metzger S, et al. Hum Genet, 2010 Oct. PMID 20697744. Plasma membrane contributes to the formation of pre-autophagosomal structures. Ravikumar B, et al. Nat Cell Biol, 2010 Aug. PMID 20639872. Replication and meta-analysis of 13,000 cases defines the risk for interleukin-23 receptor and autophagy-related 16-like 1 variants in Crohn's disease. Cotterill L, et al. Can J Gastroenterol, 2010 May. PMID 20485703. Is there a role for Crohn's disease-associated autophagy genes ATG16L1 and IRGM in formation of granulomas? Wolfkamp SC, et al. Eur J Gastroenterol Hepatol, 2010 Aug. PMID 20395867. NOD2/CARD15, ATG16L1 and IL23R gene polymorphisms and childhood-onset of Crohn's disease. Gazouli M, et al. World J Gastroenterol, 2010 Apr 14. PMID 20380008.

membrane targeting of ATG5 Localizes also
to discrete punctae along the ciliary
axoneme
{ECO:0000250|UniProtKB:Q8C0J2}

ATG16L1 Antibody - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ATG16L1 Antibody - Citations

- [Autophagy in cancer associated fibroblasts promotes tumor cell survival: Role of hypoxia, HIF1 induction and NFkB activation in the tumor stromal microenvironment.](#)