

STAT3 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8415b

Specification

STAT3 Antibody - Product Information

Application	IHC, WB,E
Primary Accession	P40763
Other Accession	P52631 , Q19S50 , P42227
Reactivity	Human
Predicted	Mouse, Pig, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a, κ
Calculated MW	88068

STAT3 Antibody - Additional Information

Gene ID 6774

Other Names

Signal transducer and activator of transcription 3, Acute-phase response factor, STAT3, APRF

Target/Specificity

This STAT3 antibody is generated from a mouse immunized with a recombinant protein from human.

Dilution

IHC~~1:5000
WB~~1:1000

Format


Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

STAT3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Immunohistochemical analysis of paraffin-embedded Human Breast cancer section using Pink1 (Cat#AM8415b). AM8415b was diluted at 1:5000 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Western blot analysis of lysates from CEM, HeLa cell line (from left to right), using Stat3 Antibody (Cat. #AM8415b). AM8415b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L (HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.

STAT3 Antibody - Background

STAT3 Antibody - Protein Information**Name** STAT3 ([HGNC:11364](#))**Function**

Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF, LEP and other growth factors (PubMed: [10688651](http://www.uniprot.org/citations/10688651) target="_blank">10688651, PubMed: [12359225](http://www.uniprot.org/citations/12359225) target="_blank">12359225, PubMed: [12873986](http://www.uniprot.org/citations/12873986) target="_blank">12873986, PubMed: [15194700](http://www.uniprot.org/citations/15194700) target="_blank">15194700, PubMed: [17344214](http://www.uniprot.org/citations/17344214) target="_blank">17344214, PubMed: [18242580](http://www.uniprot.org/citations/18242580) target="_blank">18242580, PubMed: [23084476](http://www.uniprot.org/citations/23084476) target="_blank">23084476). Once activated, recruits coactivators, such as NCOA1 or MED1, to the promoter region of the target gene (PubMed: [17344214](http://www.uniprot.org/citations/17344214) target="_blank">17344214). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed: [12873986](http://www.uniprot.org/citations/12873986) target="_blank">12873986). Upon activation of IL6ST/gp130 signaling by interleukin-6 (IL6), binds to the IL6-responsive elements identified in the promoters of various acute-phase protein genes (PubMed: [12359225](http://www.uniprot.org/citations/12359225) target="_blank">12359225). Activated by IL31 through IL31RA (PubMed: [15194700](http://www.uniprot.org/citations/15194700) target="_blank">15194700). Acts as a regulator of inflammatory response by regulating differentiation of naive CD4(+) T-cells into T-helper Th17 or regulatory T-cells (Treg): deacetylation and oxidation of lysine residues by LOXL3, leads to disrupt STAT3 dimerization and inhibit its

Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF and other growth factors. May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the interleukin-6 (IL-6)- responsive elements identified in the promoters of various acute- phase protein genes. Activated by IL31 through IL31RA. Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity. Plays an important role in host defense in methicillin-resistant S.aureus lung infection by regulating the expression of the antimicrobial lectin REG3G (By similarity).

STAT3 Antibody - References

Akira S.,et al.Cell 77:63-71(1994).
Della Pietra L.,et al.Gene 213:119-124(1998).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Della Pietra L.,et al.Submitted (OCT-1997) to the EMBL/GenBank/DBJ databases.
Zhang X.,et al.Science 267:1990-1994(1995).

transcription activity (PubMed:28065600). Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (PubMed:17344214). Mediates the effects of LEP on melanocortin production, body energy homeostasis and lactation (By similarity). May play an apoptotic role by transactivating BIRC5 expression under LEP activation (PubMed:18242580). Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity (PubMed:23084476). Plays a crucial role in basal beta cell functions, such as regulation of insulin secretion (By similarity).

Cellular Location

Cytoplasm. Nucleus. Note=Shuttles between the nucleus and the cytoplasm. Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4. Constitutive nuclear presence is independent of tyrosine phosphorylation. Predominantly present in the cytoplasm without stimuli. Upon leukemia inhibitory factor (LIF) stimulation, accumulates in the nucleus. The complex composed of BART and ARL2 plays an important role in the nuclear translocation and retention of STAT3 Identified in a complex with LYN and PAG1

Tissue Location

Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Expressed in naive CD4(+) T cells as well as T-helper Th17, Th1 and Th2 cells (PubMed:31899195)

STAT3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)