

Hsp90 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8439b

Specification

Hsp90 Antibody - Product Information

Application	WB,E
Primary Accession	P07900
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1, λ
Calculated MW	84660

Hsp90 Antibody - Additional Information

Gene ID 3320

Other Names

Heat shock protein HSP 90-alpha, Heat shock 86 kDa, HSP 86, HSP86, Lipopolysaccharide-associated protein 2, LAP-2, LPS-associated protein 2, Renal carcinoma antigen NY-REN-38, HSP90AA1, HSP90A, HSPC1, HSPCA

Target/Specificity

This antibody is generated from a mouse immunized with a recombinant protein.

Dilution

WB~~1:1000

Format


Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Hsp90 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of lysates from HeLa, K562, HepG2, mouse NIH/3T3, A2058, T47D cell line (from left to right), using Hsp90 Antibody(Cat. #AM8439b). AM8439b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 μ g per lane.

Hsp90 Antibody - Background

Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function.

Hsp90 Antibody - References

Soeda E.,et al.Nucleic Acids Res. 17:7108-7108(1989).

Hsp90 Antibody - Protein Information**Name** HSP90AA1 ([HGNC:5253](#))**Synonyms** HSP90A, HSPC1, HSPCA**Function**

Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity which is essential for its chaperone activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function (PubMed: [11274138](http://www.uniprot.org/citations/11274138), PubMed: [15577939](http://www.uniprot.org/citations/15577939), PubMed: [15937123](http://www.uniprot.org/citations/15937123), PubMed: [27353360](http://www.uniprot.org/citations/27353360), PubMed: [29127155](http://www.uniprot.org/citations/29127155), PubMed: [12526792](http://www.uniprot.org/citations/12526792)). Engages with a range of client protein classes via its interaction with various co-chaperone proteins or complexes, that act as adapters, simultaneously able to interact with the specific client and the central chaperone itself (PubMed: [29127155](http://www.uniprot.org/citations/29127155), PubMed: [29127155](http://www.uniprot.org/citations/29127155)). Recruitment of ATP and co-chaperone followed by client protein forms a functional chaperone. After the completion of the chaperoning process, properly folded client protein and co- chaperone leave HSP90 in an ADP-bound partially open conformation and finally, ADP is released from HSP90 which acquires an open conformation for the next cycle (PubMed: [27295069](http://www.uniprot.org/citations/27295069)).

Yamazaki M.,et al.Agric. Biol. Chem. 54:3163-3170(1990).

Hickey E.,et al.Mol. Cell. Biol. 9:2615-2626(1989).

Chen B.,et al.Genomics 86:627-637(2005).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

PubMed:26991466). Plays a critical role in mitochondrial import, delivers preproteins to the mitochondrial import receptor TOMM70 (PubMed:12526792). Apart from its chaperone activity, it also plays a role in the regulation of the transcription machinery. HSP90 and its co-chaperones modulate transcription at least at three different levels (PubMed:25973397). In the first place, they alter the steady-state levels of certain transcription factors in response to various physiological cues(PubMed:25973397). Second, they modulate the activity of certain epigenetic modifiers, such as histone deacetylases or DNA methyl transferases, and thereby respond to the change in the environment (PubMed:25973397). Third, they participate in the eviction of histones from the promoter region of certain genes and thereby turn on gene expression (PubMed:25973397). Binds bacterial lipopolysaccharide (LPS) and mediates LPS-induced inflammatory response, including TNF secretion by monocytes (PubMed:11276205). Antagonizes STUB1-mediated inhibition of TGF-beta signaling via inhibition of STUB1-mediated SMAD3 ubiquitination and degradation (PubMed:24613385). Mediates the association of TOMM70 with IRF3 or TBK1 in mitochondria outer membrane which promotes host antiviral response (PubMed:20628368, PubMed:25609812).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:P07901}.

Cytoplasm
{ECO:0000250|UniProtKB:P07901}.
Melanosome. Cell membrane.
Mitochondrion. Note=Identified by mass
spectrometry in melanosome fractions from
stage I to stage IV

Hsp90 Antibody - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)