

RAB5C Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8520b

Specification

RAB5C Antibody - Product Information

Application	WB, FC,E
Primary Accession	P51148
Other Accession	Q5R7L7
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG2b,k
Calculated MW	23483

RAB5C Antibody - Additional Information

Gene ID 5878

Other Names

Ras-related protein Rab-5C, L1880, RAB5L, RAB5C, RABL

Target/Specificity

This RAB5C antibody is generated from a mouse immunized with a recombinant protein of human RAB5C.

Dilution

WB~~1:2000

FC~~1:25

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

RAB5C Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RAB5C Antibody - Protein Information



All lanes : Anti-RAB5C Antibody at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: HL-60 whole cell lysate Lane 3: U-2OS whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing U-2OS cells stained with AM8520b(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AM8520b, 1:25

Name RAB5C

Synonyms RABL

Function

Protein transport. Probably involved in vesicular traffic.

Cellular Location

Cell membrane

{ECO:0000250|UniProtKB:P20339};

Lipid-anchor

{ECO:0000250|UniProtKB:P20339};

Cytoplasmic side

{ECO:0000250|UniProtKB:P20339}. Early endosome membrane

{ECO:0000250|UniProtKB:P20339};

Lipid-anchor

{ECO:0000250|UniProtKB:P20339}.

Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

RAB5C Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OJ192088) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG2b (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

RAB5C Antibody - Background

Protein transport. Probably involved in vesicular traffic (By similarity).

RAB5C Antibody - References

Han H.J., et al. Cytogenet. Cell Genet. 73:137-139(1996).

Albertsen H.M., et al. Nat. Genet. 7:472-479(1994).

Clemens D.L., et al. Infect. Immun. 68:2671-2684(2000).

Puhl H.L. III, et al. Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.

Ota T., et al. Nat. Genet. 36:40-45(2004).