

RAB14 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8601b

Specification

RAB14 Antibody - Product Information

Application	WB,E
Primary Accession	P61106
Other Accession	Q5ZKU5 , Q91V41 , Q5R8Z8
Reactivity	Human
Predicted	Chicken, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Calculated MW	23897

RAB14 Antibody - Additional Information

Gene ID 51552

Other Names

Ras-related protein Rab-14, RAB14

Target/Specificity

This RAB14 antibody is generated from a mouse immunized with a recombinant protein of human RAB14.

Dilution

WB~~1:2000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAB14 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RAB14 Antibody - Protein Information



All lanes : Anti-RAB14 Antibody at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: Raji whole cell lysate Lane 3: human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

RAB14 Antibody - Background

Involved in membrane trafficking between the Golgi complex and endosomes during early embryonic development. Regulates the Golgi to endosome transport of FGFR-containing vesicles during early development, a key process for developing basement membrane and epiblast and primitive endoderm lineages during early postimplantation development. May act by modulating the kinesin KIF16B-cargo association to endosomes (By similarity). Regulates, together with its guanine nucleotide exchange factor DENND6A, the specific endocytic transport of ADAM10, N-cadherin/CDH2 shedding and cell-cell adhesion.

RAB14 Antibody - References

Name RAB14**Function**

Involved in membrane trafficking between the Golgi complex and endosomes during early embryonic development. Regulates the Golgi to endosome transport of FGFR-containing vesicles during early development, a key process for developing basement membrane and epiblast and primitive endoderm lineages during early postimplantation development. May act by modulating the kinesin KIF16B-cargo association to endosomes (By similarity). Regulates, together with its guanine nucleotide exchange factor DENND6A, the specific endocytic transport of ADAM10, N-cadherin/CDH2 shedding and cell-cell adhesion.

Cellular Location

Recycling endosome. Early endosome membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus, trans-Golgi network membrane; Lipid-anchor; Cytoplasmic side. Cytoplasmic vesicle, phagosome. Note=Recruited to recycling endosomes by DENND6A (PubMed:22595670). Recruited to phagosomes containing S.aureus or M.tuberculosis (PubMed:21255211).

Proikas-Cezanne T.,et al.Submitted (MAY-1999) to the EMBL/GenBank/DDBJ databases.

Ren Y.,et al.Submitted (NOV-1999) to the EMBL/GenBank/DDBJ databases.

Huang Y.-P.,et al.Submitted (FEB-2004) to the EMBL/GenBank/DDBJ databases.

Hu R.-M.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).

Bechtel S.,et al.BMC Genomics 8:399-399(2007).

RAB14 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)