

HAS1 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1433a

Specification

HAS1 Antibody - Product Information

Application	WB, IF
Primary Accession	O92839
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	65kDa KDa

Description

Hyaluronan or hyaluronic acid (HA) is a high molecular weight unbranched polysaccharide synthesized by a wide variety of organisms from bacteria to mammals, and is a constituent of the extracellular matrix. It consists of alternating glucuronic acid and N-acetylglucosamine residues that are linked by beta-1-3 and beta-1-4 glycosidic bonds. HA is synthesized by membrane-bound synthase at the inner surface of the plasma membrane, and the chains are extruded through pore-like structures into the extracellular space. It serves a variety of functions, including space filling, lubrication of joints, and provision of a matrix through which cells can migrate. HA is actively produced during wound healing and tissue repair to provide a framework for ingrowth of blood vessels and fibroblasts. Changes in the serum concentration of HA are associated with inflammatory and degenerative arthropathies such as rheumatoid arthritis. In addition, the interaction of HA with the leukocyte receptor CD44 is important in tissue-specific homing by leukocytes, and overexpression of HA receptors has been correlated with tumor metastasis. HAS1 is a member of the newly identified vertebrate gene family encoding putative hyaluronan synthases, and its amino acid sequence shows significant homology to the hasA gene product of *Streptococcus pyogenes*, a glycosaminoglycan synthetase (DG42) from *Xenopus laevis*, and a recently described murine hyaluronan synthase.




Figure 1: Western blot analysis using HAS1 mAb against human HAS1 (AA: 74-243) recombinant protein. (Expected MW is 44.4 kDa)




Figure 2: Immunofluorescence analysis of U251 cells using HAS1 mAb (green). Red: Actin filaments have been labeled with DY-554 phalloidin. Blue: DRAQ5 fluorescent DNA dye.

HAS1 Antibody - References

1. Clin Lymphoma. 2005 Mar;5(4):253-6. 2.

Immunogen

Purified recombinant fragment of human HAS1 expressed in E. Coli.

Mol Cell Biochem. 2006 Nov;292(1-2):169-78.
3. J Biol Chem. 2008 Jun 13;283(24):16781-9.

Formulation

Ascitic fluid containing 0.03% sodium azide.

HAS1 Antibody - Additional Information

Gene ID 3036

Other Names

Hyaluronan synthase 1, 2.4.1.212,
Hyaluronate synthase 1, Hyaluronic acid
synthase 1, HA synthase 1, HuHAS1, HAS1,
HAS

Dilution

WB~~1/500 - 1/2000

IF~~1/200 - 1/1000

Storage

Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C in small aliquots to prevent
freeze-thaw cycles.

Precautions

HAS1 Antibody is for research use only and
not for use in diagnostic or therapeutic
procedures.

HAS1 Antibody - Protein Information

Name HAS1

Synonyms HAS

Function

Catalyzes the addition of GlcNAc or GlcUA
monosaccharides to the nascent hyaluronan
polymer. Therefore, it is essential to
hyaluronan synthesis a major component of
most extracellular matrices that has a
structural role in tissues architectures and
regulates cell adhesion, migration and
differentiation. This is one of the isozymes
catalyzing that reaction. Also able to
catalyze the synthesis of chito-
oligosaccharide depending on the substrate
(By similarity).

Cellular Location

Membrane; Multi-pass membrane protein

Tissue Location

Widely expressed. Highly expressed in ovary followed by spleen, thymus, prostate, testes and large intestine Weakly expressed in small intestine.

HAS1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)