

**CITED2 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10174B**

**Specification**

**CITED2 Antibody (C-term) - Product Information**

Application	IF, WB, E
Primary Accession	<a href="#">Q99967</a>
Other Accession	<a href="#">Q35740</a> , <a href="#">Q0VCT9</a> , <a href="#">NP_001161860.1</a> , <a href="#">NP_001161861.1</a> , <a href="#">NP_006070.2</a>
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	28497
Antigen Region	200-226

**CITED2 Antibody (C-term) - Additional Information**

**Gene ID** 10370

**Other Names**

Cbp/p300-interacting transactivator 2,  
MSG-related protein 1, MRG-1, P35srj,  
CITED2, MRG1

**Target/Specificity**

This CITED2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 200-226 amino acids from the C-terminal region of human CITED2.

**Dilution**


IF~~1:10~50  
WB~~1:1000

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C



Confocal immunofluorescent analysis of CITED2 Antibody (C-term) (Cat. #AP10174b) with Hela cell followed by Alexa Fluor® 489-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



CITED2 Antibody (C-term) (Cat. #AP10174b) western blot analysis in mouse NIH-3T3 cell line lysates (15ug/lane). This demonstrates the CITED2 antibody detected CITED2 protein (arrow).

**CITED2 Antibody (C-term) - Background**

in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

CITED2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **CITED2 Antibody (C-term) - Protein Information**

**Name** CITED2

**Synonyms** MRG1

#### **Function**

Transcriptional coactivator of the p300/CBP-mediated transcription complex. Acts as a bridge, linking TFAP2 transcription factors and the p300/CBP transcriptional coactivator complex in order to stimulate TFAP2-mediated transcriptional activation. Positively regulates TGF-beta signaling through its association with the SMAD/p300/CBP-mediated transcriptional coactivator complex. Stimulates the peroxisome proliferator-activated receptors PPARA transcriptional activity. Enhances estrogen-dependent transactivation mediated by estrogen receptors. Acts also as a transcriptional corepressor; interferes with the binding of the transcription factors HIF1A or STAT2 and the p300/CBP transcriptional coactivator complex. Participates in sex determination and early gonad development by stimulating transcription activation of SRY. Plays a role in controlling left-right patterning during embryogenesis; potentiates transcriptional activation of NODAL-mediated gene transcription in the left lateral plate mesoderm (LPM). Plays an essential role in differentiation of the adrenal cortex from the adrenogonadal primordium (AGP); stimulates WT1-mediated transcription activation thereby up-regulating the nuclear hormone receptor NR5A1 promoter activity. Associates with chromatin to the PITX2 P1 promoter region.

#### **Cellular Location**

Nucleus Note=Colocalizes with EP300 in dot-like structures

The protein encoded by this gene inhibits transactivation of HIF1A-induced genes by competing with binding of HIF1a to p300-CH1. Mutations in this gene are a cause of cardiac septal defects. Three transcript variants encoding the same protein have been found for this gene.

#### **CITED2 Antibody (C-term) - References**

Yang, X.F., et al. Zhonghua Er Ke Za Zhi 48(4):293-296(2010)  
Sun, H.B. Ann. N. Y. Acad. Sci. 1192, 429-436 (2010) :  
Lau, W.M., et al. Int. J. Cancer 126(4):876-884(2010)  
van Agthoven, T., et al. Br. J. Cancer 101(11):1824-1832(2009)  
Ganesh, S.K., et al. Nat. Genet. 41(11):1191-1198(2009)

#### **CITED2 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)