

CCDC148 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10609c

Specification

CCDC148 Antibody (Center) - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
WB, FC,E
O8NFR7
NP_620158.3
Human, Mouse
Rabbit

Host
Clonality
Isotype
Calculated MW
Antigen Region

Rabbit
Polyclonal
Rabbit Ig
71076
279-307

CCDC148 Antibody (Center) - Additional Information

Gene ID 130940

Other Names

Coiled-coil domain-containing protein 148, CCDC148

Target/Specificity

This CCDC148 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 279-307 amino acids from the Central region of human CCDC148.

Dilution

WB~~1:1000 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CCDC148 Antibody (Center) is for research



CCDC148 Antibody (Center) (Cat. #AP10609c) western blot analysis in A2058 cell line lysates (35ug/lane). This demonstrates the CCDC148 antibody detected the CCDC148 protein (arrow).



CCDC148 Antibody (Center) (Cat. #AP10609c) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the CCDC148 antibody detected the CCDC148 protein (arrow).



use only and not for use in diagnostic or therapeutic procedures.


CCDC148 Antibody (Center) - Protein Information

Name CCDC148

CCDC148 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



CCDC148 Antibody (Center) (Cat. #AP10609c) flow cytometric analysis of A2058 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CCDC148 Antibody (Center) - Background

There are three different isoforms.