

JMJD2B Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11950a

Specification

JMJD2B Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O94953
Other Accession	NP_055830
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	121897
Antigen Region	428-459

JMJD2B Antibody (N-term) - Additional Information

Gene ID 23030

Other Names

Lysine-specific demethylase 4B, 11411-, JmjC domain-containing histone demethylation protein 3B, Jumonji domain-containing protein 2B, KDM4B, JHDM3B, JMJD2B, KIAA0876

Target/Specificity

This JMJD2B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 428-459 amino acids from the N-terminal region of human JMJD2B.

Dilution


WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



JMJD2B Antibody (N-term) (Cat. #AP11950a) western blot analysis in A375 cell line lysates (35ug/lane). This demonstrates the JMJD2B antibody detected the JMJD2B protein (arrow).

JMJD2B Antibody (N-term) - Background

KDM4B is a histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a role in histone code. It does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys-20' and only able to demethylate trimethylated H3 'Lys-9', with a weaker activity than KDM4A, KDM4C and KDM4D. Demethylation of Lys residue generates formaldehyde and succinate.

JMJD2B Antibody (N-term) - References

- Yang, J., et al. Cancer Res. 70(16):6456-6466(2010)
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Beyer, S., et al. J. Biol. Chem. 283(52):36542-36552(2008)
Pollard, P.J., et al. Biochem. J. 416(3):387-394(2008)
Katoh, Y., et al. Int. J. Mol. Med. 20(2):269-273(2007)

Precautions

JMJD2B Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

JMJD2B Antibody (N-term) - Protein Information

Name KDM4B

Synonyms JHDM3B, JMJD2B, KIAA0876

Function

Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys- 20'. Only able to demethylate trimethylated H3 'Lys-9', with a weaker activity than KDM4A, KDM4C and KDM4D. Demethylation of Lys residue generates formaldehyde and succinate.

Cellular Location

Nucleus

{ECO:0000255|PROSITE-ProRule:PRU00537,
ECO:0000269|PubMed:15927959}

JMJD2B Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)