

## **PSCD2 Antibody (N-term)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12239A

## **Specification**

#### PSCD2 Antibody (N-term) - Product Information

Application WB,E
Primary Accession 099418

Other Accession P63035, P63034,

Q2KI41,

NP 004219.3

Reactivity
Predicted
Host
Clonality
Isotype
Calculated MW
Antigen Region

Mouse
Bovine, Rat
Rabbit
Polyclonal
Rabbit Ig
46546
36-64

PSCD2 Antibody (N-term) - Additional Information

## **Gene ID 9266**

#### **Other Names**

Cytohesin-2, ARF exchange factor, ARF nucleotide-binding site opener, Protein ARNO, PH, SEC7 and coiled-coil domain-containing protein 2, CYTH2, ARNO, PSCD2, PSCD2L

# **Target/Specificity**

This PSCD2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 36-64 amino acids from the N-terminal region of human PSCD2.

#### **Dilution**


WB~~1:1000

## **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

## Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



PSCD2 Antibody (N-term) (Cat. #AP12239a) western blot analysis in mouse kidney tissue lysates (35ug/lane). This demonstrates the PSCD2 antibody detected the PSCD2 protein (arrow).

## PSCD2 Antibody (N-term) - Background

The protein encoded by this gene is a member of the PSCD

family. Members of this family have identical structural

organization that consists of an N-terminal coiled-coil motif, a

central Sec7 domain, and a C-terminal pleckstrin homology (PH)

domain. The coiled-coil motif is involved in homodimerization, the

Sec7 domain contains guanine-nucleotide exchange protein (GEP)

activity, and the PH domain interacts with phospholipids and is


responsible for association of PSCDs with membranes. Members of

this family appear to mediate the regulation of protein sorting and

membrane trafficking. The encoded protein exhibits GEP activity in

vitro with ARF1, ARF3, and ARF6 and is 83% homologous to CYTH1. Two

transcript variants encoding different isoforms





Tel: 858.875.1900 Fax: 858.622.0609

cycles.

#### **Precautions**

PSCD2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PSCD2 Antibody (N-term) - Protein Information

Name CYTH2 (HGNC:9502)

Synonyms ARNO, PSCD2, PSCD2L

#### **Function**

Acts as a guanine-nucleotide exchange factor (GEF). Promotes quanine-nucleotide exchange on ARF1, ARF3 and ARF6. Activates ARF factors through replacement of GDP with GTP (By similarity). The cell membrane form, in association with ARL4 proteins, recruits ARF6 to the plasma membrane (PubMed: <a href="http://www.u niprot.org/citations/17398095" target=" blank">17398095</a>). Involved in neurite growth (By similarity).

## **Cellular Location**

Cell membrane; Peripheral membrane protein. Cytoplasm. Cell projection {ECO:0000250|UniProtKB:P63034}. Cell projection, growth cone {ECO:0000250|UniProtKB:P63034}. Cell junction, tight junction {ECO:0000250|UniProtKB:P63034}. Cell junction, adherens junction {ECO:0000250|UniProtKB:P63034}. Note=Both isoform 1 and isoform 2 are recruited to the cell membrane through its association with ARL4A, ARL4C and ARL4D. They require also interaction with phosphoinositides for targeting to plasma membrane (PubMed:17398095). In differentiating neuroblastoma cells, colocalizes with CCDC120 in both neurite shaft and growth cone areas. {ECO:0000250|UniProtKB:P63034, ECO:0000269|PubMed:17398095}

**Tissue Location** Widely expressed..


# PSCD2 Antibody (N-term) - Protocols

Provided below are standard protocols that you

have been found for this gene.

# **PSCD2** Antibody (N-term) - References

Merkulova, M., et al. Biochim. Biophys. Acta 1797(8):1398-1409(2010) Torii, T., et al. J. Biol. Chem. 285(31):24270-24281(2010) White, D.T., et al. Mol. Biol. Cell 21(4):562-571(2010) Bouschet, T., et al. J. Cell. Sci. 120 (PT 15), 2489-2497 (2007): Wu, C., et al. Proteomics 7(11):1775-1785(2007)





may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture