

**IRAK4 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP12875A**

**Specification**

**IRAK4 Antibody (N-term) - Product Information**

Application	<b>WB,E</b>
Primary Accession	<a href="#">Q9NWZ3</a>
Other Accession	<a href="#">Q8R4K2</a> , <a href="#">Q1RMT8</a> , <a href="#">NP_001107654.1</a> , <a href="#">NP_057207.2</a>
Reactivity	<b>Human</b>
Predicted	<b>Bovine, Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Antigen Region	<b>25-52</b>

**IRAK4 Antibody (N-term) - Additional Information**

**Gene ID** 51135

**Other Names**

Interleukin-1 receptor-associated kinase 4,  
IRAK-4, Renal carcinoma antigen  
NY-REN-64, IRAK4

**Target/Specificity**

This IRAK4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 25-52 amino acids from the N-terminal region of human IRAK4.

**Dilution**

WB ~ ~ 1:1000


**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**



IRAK4 Antibody (N-term) (Cat. #AP12875a) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the IRAK4 antibody detected the IRAK4 protein (arrow).



Anti-IRAK4 Antibody (N-term) at 1:1000 dilution + K562 whole cell lysate  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 52 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

IRAK4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **IRAK4 Antibody (N-term) - Protein Information**

##### **Name** IRAK4

##### **Function**

Serine/threonine-protein kinase that plays a critical role in initiating innate immune response against foreign pathogens. Involved in Toll-like receptor (TLR) and IL-1R signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/17878374" target="\_blank">17878374</a>). Is rapidly recruited by MYD88 to the receptor-signaling complex upon TLR activation to form the Myddosome together with IRAK2. Phosphorylates initially IRAK1, thus stimulating the kinase activity and intensive autophosphorylation of IRAK1. Phosphorylates E3 ubiquitin ligases Pellino proteins (PELI1, PELI2 and PELI3) to promote pellino-mediated polyubiquitination of IRAK1. Then, the ubiquitin-binding domain of IKBKG/NEMO binds to polyubiquitinated IRAK1 bringing together the IRAK1-MAP3K7/TAK1-TRAF6 complex and the NEMO-IKKA-IKKB complex. In turn, MAP3K7/TAK1 activates IKKs (CHUK/IKKA and IKBKB/IKKB) leading to NF-kappa-B nuclear translocation and activation. Alternatively, phosphorylates TIRAP to promote its ubiquitination and subsequent degradation. Phosphorylates NCF1 and regulates NADPH oxidase activation after LPS stimulation suggesting a similar mechanism during microbial infections.

##### **Cellular Location**

Cytoplasm.

#### **IRAK4 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

#### **IRAK4 Antibody (N-term) - Background**

This gene encodes a kinase that activates NF-kappaB in both the Toll-like receptor (TLR) and T-cell receptor (TCR) signaling pathways. The protein is essential for most innate immune responses. Mutations in this gene result in IRAK4 deficiency and recurrent invasive pneumococcal disease. Multiple transcript variants encoding different isoforms have been found for this gene.

#### **IRAK4 Antibody (N-term) - References**

- Silva, L.K., et al. Eur. J. Hum. Genet. 18(11):1221-1227(2010)  
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)  
McDonald, D.R., et al. J. Allergy Clin. Immunol. 126(2):332-337(2010)  
Schuurhof, A., et al. Pediatr. Pulmonol. 45(6):608-613(2010)  
Wang, Z., et al. Structure 14(12):1835-1844(2006)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**IRAK4 Antibody (N-term) - Citations**

- [The anti-inflammatory effect and potential mechanism of cardamonin in DSS-induced colitis.](#)