

SERPINE1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12919C

Specification

SERPINE1 Antibody (Center) - Product Information

Application WB, FC,E
Primary Accession
Other Accession NP_000593.1,
NP_001158885.1

Reactivity
Host
Clonality
Isotype
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
188-216

SERPINE1 Antibody (Center) - Additional Information

Gene ID 5054

Other Names

Plasminogen activator inhibitor 1, PAI, PAI-1, Endothelial plasminogen activator inhibitor, Serpin E1, SERPINE1, PAI1, PLANH1

Target/Specificity

This SERPINE1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 188-216 amino acids from the Central region of human SERPINE1.

Dilution


WB~~1:1000 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western blot analysis of SERPINE1 (arrow) using rabbit polyclonal SERPINE1 Antibody (Center) (Cat. #AP12919c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the SERPINE1 gene.



SERPINE1 Antibody (Center) (Cat. #AP12919c) flow cytometric analysis of U251 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Precautions

SERPINE1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

SERPINE1 Antibody (Center) - Protein Information

Name SERPINE1

Synonyms PAI1, PLANH1

Function

Serine protease inhibitor. Inhibits TMPRSS7 (PubMed:<a href="http://www.uniprot.org/c itations/15853774"

target="_blank">15853774). Is a primary inhibitor of tissue-type plasminogen activator (PLAT) and urokinase-type plasminogen activator (PLAU). As PLAT inhibitor, it is required for fibrinolysis down-regulation and is responsible for the controlled degradation of blood clots (PubMed:<a href="http://www.uniprot.org/c itations/8481516"

target="_blank">8481516,

PubMed: <a href="http://www.uniprot.org/ci tations/9207454"

target="_blank">9207454,

PubMed:<a href="http://www.uniprot.org/ci tations/17912461"

target="_blank">17912461). As PLAU inhibitor, it is involved in the regulation of cell adhesion and spreading (PubMed:<a hr ef="http://www.uniprot.org/citations/9175705" target="_blank">9175705). Acts as a regulator of cell migration, independently of its role as protease inhibitor (PubMed:http://www.unip

rot.org/citations/15001579" target="_blank">15001579,

PubMed:<a href="http://www.uniprot.org/ci tations/9168821"

target="_blank">9168821). It is required for stimulation of keratinocyte migration during cutaneous injury repair (PubMed:<a href="http://www.uniprot.org/citations/18386027"

target="_blank">18386027). It is involved in cellular and replicative senescence (PubMed:16862142). Plays a role in alveolar type 2 cells senescence in

role in alveolar type 2 cells senescence in the lung (By similarity). Is involved in the regulation of cementogenic differentiation of periodontal ligament stem cells, and

SERPINE1 Antibody (Center) - Background

This gene encodes a member of the serine proteinase

inhibitor (serpin) superfamily. This member is the principal

inhibitor of tissue plasminogen activator (tPA) and urokinase

(uPA), and hence is an inhibitor of fibrinolysis. Defects in this

gene are the cause of plasminogen activator inhibitor-1 deficiency

(PAI-1 deficiency), and high concentrations of the gene product are

associated with thrombophilia. Alternatively spliced transcript

variants encoding different isoforms have been found for this gene.

SERPINE1 Antibody (Center) - References

Ma, Z., et al. Biochem. Biophys. Res. Commun. 400(4):569-574(2010)


Ince, D.A., et al. Genet Test Mol Biomarkers 14(5):643-647(2010)

Bern, M.M., et al. Clin. Appl. Thromb. Hemost. 16(5):574-578(2010)

Markl, B., et al. J Surg Oncol

102(3):235-241(2010)

de Haas, E.C., et al. Cancer (2010) In press :





regulates odontoblast differentiation and dentin formation during odontogenesis (PubMed:25808697, PubMed:27046084).

Cellular Location Secreted.

Tissue Location

Expressed in endothelial cells (PubMed:2430793, PubMed:3097076). Found in plasma, platelets, and hepatoma and fibrosarcoma cells.

SERPINE1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture