

MPO Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13056B

Specification

MPO Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P05164
Other Accession	NP_000241.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	83869
Antigen Region	548-575

MPO Antibody (C-term) - Additional Information

Gene ID 4353

Other Names

Myeloperoxidase, MPO, Myeloperoxidase, 89 kDa myeloperoxidase, 84 kDa myeloperoxidase, Myeloperoxidase light chain, Myeloperoxidase heavy chain, MPO

Target/Specificity

This MPO antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 548-575 amino acids from the C-terminal region of human MPO.

Dilution

WB~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MPO Antibody (C-term) is for research use



MPO Antibody (C-term) (Cat. #AP13056b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the MPO antibody detected the MPO protein (arrow).

MPO Antibody (C-term) - Background

Myeloperoxidase (MPO) is a heme protein synthesized during myeloid differentiation that constitutes the major component of neutrophil azurophilic granules. Produced as a single chain precursor, myeloperoxidase is subsequently cleaved into a light and heavy chain. The mature myeloperoxidase is a tetramer composed of 2 light chains and 2 heavy chains. This enzyme produces hypohalous acids central to the microbicidal activity of neutrophils. [provided by RefSeq].

MPO Antibody (C-term) - References

Banerjee, M., et al. Toxicol. Appl. Pharmacol. 249(1):47-54(2010)
Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)
Nahon, P., et al. Antioxid. Redox Signal. (2010)

only and not for use in diagnostic or therapeutic procedures.

In press :
Wang, Y., et al. J. Huazhong Univ. Sci. Technol. Med. Sci. 30(4):437-442(2010)
Hua, F., et al. Zhongguo Fei Ai Za Zhi 13(2):122-127(2010)

MPO Antibody (C-term) - Protein Information

Name MPO

Function

Part of the host defense system of polymorphonuclear leukocytes. It is responsible for microbicidal activity against a wide range of organisms. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity.

Cellular Location

Lysosome.

MPO Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)