

MPP3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13068b

Specification

MPP3 Antibody (C-term) - Product Information

Application	IF, WB, FC, E
Primary Accession	Q13368
Other Accession	NP_001923.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	66152
Antigen Region	406-434

MPP3 Antibody (C-term) - Additional Information

Gene ID 4356

Other Names

MAGUK p55 subfamily member 3, Discs large homolog 3, Protein MPP3, MPP3, DLG3

Target/Specificity

This MPP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 406-434 amino acids from the C-terminal region of human MPP3.

Dilution

IF~~1:10~50

WB~~1:1000

FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MPP3 Antibody (C-term) is for research use



Confocal immunofluorescent analysis of MPP3 Antibody (C-term)(Cat#AP13068b) with MDA-MB231 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green).DAPI was used to stain the cell nuclear (blue).



MPP3 Antibody (C-term) (Cat. #AP13068b) western blot analysis in K562,NCI-H292 cell line lysates (35ug/lane).This demonstrates the MPP3 antibody detected the MPP3 protein (arrow).

only and not for use in diagnostic or therapeutic procedures.

MPP3 Antibody (C-term) - Protein Information


Name MPP3

Synonyms DLG3


MPP3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Anti-MPP3 Antibody (C-term) at 1:1000 dilution + HepG2 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 66 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



MPP3 Antibody (C-term) (Cat. #AP13068b) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.

MPP3 Antibody (C-term) - Background

This gene product is a member of a family of membrane-associated proteins termed MAGUKs (membrane-associated guanylate kinase homologs). MAGUKs interact with the cytoskeleton

and regulate cell proliferation, signaling pathways, and intracellular junctions. This protein contains a conserved sequence, called the SH3 (src homology 3) motif, found in several other proteins that associate with the cytoskeleton and are suspected to play important roles in signal transduction. Alternatively spliced transcript variants have been identified. One transcript variant is experimentally supported, but it doesn't encode a protein.

MPP3 Antibody (C-term) - References

Qin, L., et al. Zhonghua Gan Zang Bing Za Zhi 15(7):509-512(2007)
Kantardzhieva, A., et al. FEBS J. 273(6):1152-1165(2006)
Stelzl, U., et al. Cell 122(6):957-968(2005)
Fukuhara, H., et al. Oncogene 22(40):6160-6165(2003)
Shin, B.K., et al. J. Biol. Chem. 278(9):7607-7616(2003)