

CAPN8 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13161c

Specification

CAPN8 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	A6NHC0
Other Accession	NP_001137434.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	79144
Antigen Region	290-318

CAPN8 Antibody (Center) - Additional Information

Gene ID 388743

Other Names

Calpain-8, New calpain 2, nCL-2,
Stomach-specific M-type calpain, CAPN8,
NCL2

Target/Specificity

This CAPN8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 290-318 amino acids from the Central region of human CAPN8.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CAPN8 Antibody (Center) is for research use



CAPN8 Antibody (Center) (Cat. #AP13161c) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the CAPN8 antibody detected the CAPN8 protein (arrow).



CAPN8 Antibody (Center) (Cat. #AP13161c) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CAPN8 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

only and not for use in diagnostic or therapeutic procedures.

CAPN8 Antibody (Center) - Protein Information

Name CAPN8

Synonyms NCL2

Function

Calcium-regulated non-lysosomal thiol-protease. Involved in membrane trafficking in the gastric surface mucus cells (pit cells) and may involve the membrane trafficking of mucus cells via interactions with coat protein. Proteolytically cleaves the beta-subunit of coatomer complex (By similarity).

Cellular Location

Cytoplasm. Golgi apparatus

Tissue Location

Stomach.

CAPN8 Antibody (Center) - Background

Calcium-regulated non-lysosomal thiol-protease. Involved in membrane trafficking in the gastric surface mucus cells (pit cells) and may involve the membrane trafficking of mucus cells via interactions with coat protein. Proteolytically cleaves the beta-subunit of coatomer complex (By similarity).

CAPN8 Antibody (Center) - References

Hata, S., et al. J. Biol. Chem. 282(38):27847-27856(2007)
Oh, J.H., et al. Mamm. Genome 16(12):942-954(2005)
Hata, S., et al. J. Mol. Evol. 53(3):191-203(2001)
Huang, Y., et al. Trends Mol Med 7(8):355-362(2001)
Harrington, J.J., et al. Nat. Biotechnol. 19(5):440-445(2001)

CAPN8 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)