

TUBB8 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13248a

Specification

TUBB8 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q3ZCM7
Other Accession	NP_817124.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	49776
Antigen Region	34-62

TUBB8 Antibody (N-term) - Additional Information

Gene ID 347688

Other Names

Tubulin beta-8 chain, TUBB8

Target/Specificity

This TUBB8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 34-62 amino acids from the N-terminal region of human TUBB8.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TUBB8 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



TUBB8 Antibody (N-term) (Cat. #AP13248a) western blot analysis in 293,A549,HepG2,K562 cell line lysates (35ug/lane).This demonstrates the TUBB8 antibody detected the TUBB8 protein (arrow).

TUBB8 Antibody (N-term) - Background

Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain (By similarity).

TUBB8 Antibody (N-term) - References

van Geel, M., et al. Am. J. Hum. Genet. 70(1):269-278(2002)
van Geel, M., et al. Cytogenet. Cell Genet. 88 (3-4), 316-321 (2000) :

TUBB8 Antibody (N-term) - Protein Information**Name** TUBB8 ([HGNC:20773](#))**Function**

Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain (By similarity). TUBB8 has a key role in meiotic spindle assembly and oocyte maturation (PubMed:26789871).

Cellular Location

Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, spindle

Tissue Location

Expressed at a high level in oocytes, at different stages of development.

TUBB8 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)