

PCDH18 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13587B

Specification

PCDH18 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9HCL0
Other Accession	NP_061908.1
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	126149
Antigen Region	970-999

PCDH18 Antibody (C-term) - Additional Information

Gene ID 54510

Other Names

Protocadherin-18, PCDH18, KIAA1562

Target/Specificity

This PCDH18 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 970-999 amino acids from the C-terminal region of human PCDH18.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PCDH18 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



PCDH18 Antibody (C-term) - Background

This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily. This gene encodes a protein which contains 6 extracellular cadherin domains, a transmembrane domain and a cytoplasmic tail differing from those of the classical cadherins. Although its specific function is undetermined, the cadherin-related neuronal receptor is thought to play a role in the establishment and function of specific cell-cell connections in the brain.

PCDH18 Antibody (C-term) - References

Homayouni, R., et al. Biochem. Biophys. Res. Commun. 289(2):539-547(2001)

PCDH18 Antibody (C-term) - Protein Information**Name** PCDH18**Synonyms** KIAA1562**Function**

Potential calcium-dependent cell-adhesion protein.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed in all tissues, with highest expression in lung and ovary.

Wolverton, T., et al. Genomics 76 (1-3), 66-72 (2001) :

Suzuki, S.T. Exp. Cell Res. 261(1):13-18(2000)

Nollet, F., et al. J. Mol. Biol. 299(3):551-572(2000)

Yagi, T., et al. Genes Dev. 14(10):1169-1180(2000)

PCDH18 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)