

HIST2H2AA3 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13700a

Specification

HIST2H2AA3 Antibody (N-term) - Product Information

Application	WB, E
Primary Accession	Q6FI13
Other Accession	P09588 , P27661 , P16104 , A9UMV8 , Q8R1M2 , Q4R3X5 , Q9BTM1 , P70082 , Q3ZBX9 , P02263 , Q4FZT6 , Q8BFU2 , Q7L7L0 , P35062 , Q64523 , Q16777 , A1A4R1 , Q64522 , Q8IUE6 , P0CC09 , Q6GSS7 , P02262 , P0C169 , Q93077 , Q96QV6 , NP_001035807.1 , NP_003507.1
Reactivity Predicted	Human Rat, Mouse, Bovine, Chicken, Monkey, C.Elegans
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	14095
Antigen Region	2-30

HIST2H2AA3 Antibody (N-term) - Additional Information


Gene ID 723790;8337

Other Names

Histone H2A type 2-A, Histone H2A2,
Histone H2A/o, HIST2H2AA3, H2AFO,
HIST2H2AA

Target/Specificity

This HIST2H2AA3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 2-30 amino acids from the N-terminal region of human HIST2H2AA3.



HIST2H2AA3 Antibody (N-term) (Cat. #AP13700a) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the HIST2H2AA3 antibody detected the HIST2H2AA3 protein (arrow).

HIST2H2AA3 Antibody (N-term) - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HIST2H2AA3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

four histone genes in the cluster that are duplicated; this record represents the centromeric copy.

HIST2H2AA3 Antibody (N-term) - References

Bergink, S., et al. Genes Dev. 20(10):1343-1352(2006)
Bonenfant, D., et al. Mol. Cell Proteomics 5(3):541-552(2006)
Boyne, M.T. II, et al. J. Proteome Res. 5(2):248-253(2006)
Cao, R., et al. Mol. Cell 20(6):845-854(2005)
Hagiwara, T., et al. Biochemistry 44(15):5827-5834(2005)

HIST2H2AA3 Antibody (N-term) - Protein Information

Name H2AC18 ([HGNC:4736](#))

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location

Nucleus. Chromosome.

HIST2H2AA3 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)