

IL21 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13840b

Specification

IL21 Antibody (C-term) - Product Information

Application WB,E Q9HBE4 Primary Accession Other Accession NP 068575.1 Reactivity Human Host Rabbit Clonality **Polyclonal** Isotype Rabbit Ig Calculated MW 18653 Antigen Region 104-132

IL21 Antibody (C-term) - Additional Information

Gene ID 59067

Other Names

Interleukin-21, IL-21, Za11, IL21

Target/Specificity

This IL21 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 104-132 amino acids from the C-terminal region of human IL21.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IL21 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



IL21 Antibody (C-term) (Cat. #AP13840b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the IL21 antibody detected the IL21 protein (arrow).



Western blot analysis of IL21 (arrow) using rabbit polyclonal IL21 Antibody (C-term) (Cat. #AP13840b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the IL21 gene.

IL21 Antibody (C-term) - Background

Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of lgG(1) and lgG(3) in B-cells (By similarity). May



IL21 Antibody (C-term) - Protein Information

Name IL21 (HGNC:6005)

Function

Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity), Implicated in the generation and maintenance of T follicular helper (Tfh) cells and the formation of germinal-centers. Together with IL6, control the early generation of Tfh cells and are critical for an effective antibody response to acute viral infection (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells (PubMed:11081504, PubMed:15178704). During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation (By similarity).

Cellular Location Secreted.

Tissue Location

Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes

IL21 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B-and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation.

IL21 Antibody (C-term) - References

Petukhova, L., et al. Nature 466(7302):113-117(2010)
Yue, F.Y., et al. J. Immunol. 185(1):498-506(2010)
Pallone, F., et al. Expert Rev Clin Immunol 6(4):537-541(2010)
Schuurhof, A., et al. Pediatr. Pulmonol. 45(6):608-613(2010)
Hollis-Moffatt, J.E., et al. Arthritis Res. Ther. 12 (3), R116 (2010):