

OCLN Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13901b

Specification

OCLN Antibody (C-term) - Product Information

Application WB,E
Primary Accession Other Accession Q61146,
NP 002529.1

Reactivity
Predicted
Host
Clonality
Isotype
Calculated MW
Antigen Region

Human
Mouse
Rabbit
Polyclonal
Rabbit Ig
59144
446-474

OCLN Antibody (C-term) - Additional Information

Gene ID 100506658

Other Names Occludin, OCLN

Target/Specificity

This OCLN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 446-474 amino acids from the C-terminal region of human OCLN.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OCLN Antibody (C-term) is for research use only and not for use in diagnostic or




Western blot analysis of lysate from MCF-7 cell line, using OCLN Antibody (C-term)(Cat. #AP13901b). AP13901b was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.

OCLN Antibody (C-term) - Background

This gene encodes an integral membrane protein which is located at tight junctions. This protein may be involved in the formation and maintenance of the tight junction. The possibility of several alternatively spliced products has been suggested but the full nature of these products has not been described. [provided by RefSeq].

OCLN Antibody (C-term) - References

Liu, S., et al. Virology 407(1):160-170(2010) Michta, M.L., et al. J. Virol. 84(22):11696-11708(2010) O'Driscoll, M.C., et al. Am. J. Hum. Genet. 87(3):354-364(2010)



Tel: 858.875.1900 Fax: 858.622.0609



therapeutic procedures.

OCLN Antibody (C-term) - Protein Information

Name OCLN

Function

May play a role in the formation and regulation of the tight junction (TJ) paracellular permeability barrier. It is able to induce adhesion when expressed in cells lacking tight junctions.

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell junction, tight junction

Tissue Location

Localized at tight junctions of both epithelial and endothelial cells. Highly expressed in kidney. Not detected in testis

OCLN Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cvtometv
- Cell Culture

Van Itallie, C.M., et al. J. Cell. Sci. 123 (PT 16), 2844-2852 (2010): Raikwar, N.S., et al. Am. J. Physiol. Renal Physiol. 299 (2), F436-F444 (2010):