

BBC3 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14100c

Specification

BBC3 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q9BXH1
Other Accession	NP_055232.1 , NP_001120714.1 , NP_001120713.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	20532
Antigen Region	109-137

BBC3 Antibody (Center) - Additional Information

Gene ID 27113

Other Names

Bcl-2-binding component 3, JFY-1, p53 up-regulated modulator of apoptosis, BBC3, PUMA

Target/Specificity

This BBC3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 109-137 amino acids from the Central region of human BBC3.

Dilution


WB~~1:1000
IHC-P~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



BBC3 Antibody (Center) (Cat. #AP14100c) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the BBC3 antibody detected the BBC3 protein (arrow).



BBC3 Antibody (Center) (Cat. #AP14100c) immunohistochemistry analysis in formalin fixed and paraffin embedded human cervix tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of BBC3 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Precautions

BBC3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

BBC3 Antibody (Center) - Protein Information

Name BBC3

Synonyms PUMA

Function

Essential mediator of p53/TP53-dependent and p53/TP53- independent apoptosis (PubMed:<a href="http://www.uniprot.org/citations/11463391"

target="_blank">11463391).

Functions by promoting partial unfolding of BCL2L1 and dissociation of BCL2L1 from p53/TP53. Regulates ER stress-induced neuronal apoptosis (PubMed:23340338).

Cellular Location

Mitochondrion Note=Localized to the mitochondria in order to induce cytochrome c release

Tissue Location

Ubiquitously expressed.

BBC3 Antibody (Center) - Background

BBC3 is an essential mediator of p53-dependent and p53-independent apoptosis.

BBC3 Antibody (Center) - References

Cazanave, S.C., et al. Am. J. Physiol. Gastrointest. Liver Physiol. 299 (1), G236-G243 (2010) :

Gurzov, E.N., et al. J. Biol. Chem. 285(26):19910-19920(2010)

Hemmati, P.G., et al. J. Mol. Med. 88(6):609-622(2010)

Gomes, N.P., et al. Genes Dev. 24(10):1022-1034(2010)

Liu, F.T., et al. Biochem. Biophys. Res. Commun. 310(3):956-962(2003)

BBC3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)