

RCAN3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14397b

Specification

RCAN3 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9UKA8
Other Accession	NP_038469.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	27492
Antigen Region	194-222

RCAN3 Antibody (C-term) - Additional Information

Gene ID 11123

Other Names

Calcipressin-3, Down syndrome candidate region 1-like protein 2, Myocyte-enriched calcineurin-interacting protein 3, MCIP3, Regulator of calcineurin 3, RCAN3, DSCR1L2

Target/Specificity

This RCAN3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 194-222 amino acids from the C-terminal region of human RCAN3.

Dilution

WB~~1:1000
IHC-P~~1:10~50


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



RCAN3 Antibody (C-term) (Cat. #AP14397b) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the RCAN3 antibody detected the RCAN3 protein (arrow).



RCAN3 Antibody (C-term) (AP14397b) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of RCAN3 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

RCAN3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RCAN3 Antibody (C-term) - Protein Information

Name RCAN3

Synonyms DSCR1L2

Function

Inhibits calcineurin-dependent transcriptional responses by binding to the catalytic domain of calcineurin A. Could play a role during central nervous system development (By similarity).

Tissue Location

Highest expression in heart, skeletal muscle kidney, liver and peripheral blood leukocytes. Lower expression in all other tissues

RCAN3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RCAN3 Antibody (C-term) - Background

RCAN3 inhibits calcineurin-dependent transcriptional responses by binding to the catalytic domain of calcineurin A. Could play a role during central nervous system development (By similarity).

RCAN3 Antibody (C-term) - References

Sales, K.J., et al. *Biochim. Biophys. Acta* 1793(12):1917-1928(2009)
Enjuanes, A., et al. *Cancer Res.* 68(24):10178-10186(2008)
Facchin, F., et al. *Gene* 407 (1-2), 159-168 (2008) :
Mulero, M.C., et al. *Biochim. Biophys. Acta* 1773(3):330-341(2007)
Olsen, J.V., et al. *Cell* 127(3):635-648(2006)