

**IDE Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1455c**

**Specification**

**IDE Antibody (Center) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">P14735</a>
Other Accession	<a href="#">P35559</a> , <a href="#">Q9JHR7</a> , <a href="#">Q24K02</a>
Reactivity	Human, Mouse
Predicted	Bovine, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	406-435

**IDE Antibody (Center) - Additional Information**

**Gene ID 3416**

**Other Names**

Insulin-degrading enzyme, Abeta-degrading protease, Insulin protease, Insulinase, Insulysin, IDE

**Target/Specificity**

This IDE antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 406-435 amino acids from the Central region of human IDE.

**Dilution**

WB~~1:2000  
 IHC-P~~1:10~50

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

IDE Antibody (Center) is for research use



All lanes : Anti-IDE Antibody (Center) at 1:2000 dilution Lane 1: Hela whole cell lysate  
 Lane 2: HepG2 whole cell lysate Lane 3: A375 whole cell lysate Lane 4: K562 whole cell lysate Lane 5: mouse liver lysate  
 Lysates/proteins at 20 µg per lane.  
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.  
 Predicted band size : 118 kDa  
 Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of lysate from K562

only and not for use in diagnostic or therapeutic procedures.

#### IDE Antibody (Center) - Protein Information



##### Name IDE

{ECO:0000303|PubMed:20364150,  
 ECO:0000312|HGNC:HGNC:5381}

##### Function

Plays a role in the cellular breakdown of insulin, APP peptides, IAPP peptides, natriuretic peptides, glucagon, bradykinin, kallidin, and other peptides, and thereby plays a role in intercellular peptide signaling (PubMed:<a href="http://www.uniprot.org/citations/2293021" target="\_blank">2293021</a>, PubMed:<a href="http://www.uniprot.org/citations/10684867" target="\_blank">10684867</a>, PubMed:<a href="http://www.uniprot.org/citations/26968463" target="\_blank">26968463</a>, PubMed:<a href="http://www.uniprot.org/citations/17051221" target="\_blank">17051221</a>, PubMed:<a href="http://www.uniprot.org/citations/17613531" target="\_blank">17613531</a>, PubMed:<a href="http://www.uniprot.org/citations/18986166" target="\_blank">18986166</a>, PubMed:<a href="http://www.uniprot.org/citations/19321446" target="\_blank">19321446</a>, PubMed:<a href="http://www.uniprot.org/citations/23922390" target="\_blank">23922390</a>, PubMed:<a href="http://www.uniprot.org/citations/24847884" target="\_blank">24847884</a>, PubMed:<a href="http://www.uniprot.org/citations/26394692" target="\_blank">26394692</a>, PubMed:<a href="http://www.uniprot.org/citations/29596046" target="\_blank">29596046</a>, PubMed:<a href="http://www.uniprot.org/citations/21098034" target="\_blank">21098034</a>). Substrate binding induces important conformation changes, making it possible to bind and degrade larger substrates, such as insulin (PubMed:<a href="http://www.uniprot.org/citations/23922390" target="\_blank">23922390</a>).

cell line, using IDE Antibody (Center)(Cat. #AP1455c). AP1455c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.




target="\_blank">>23922390</a>,  
PubMed:<a href="http://www.uniprot.org/citations/26394692"  
target="\_blank">>26394692</a>,  
PubMed:<a href="http://www.uniprot.org/citations/29596046"  
target="\_blank">>29596046</a>).  
Contributes to the regulation of peptide hormone signaling cascades and regulation of blood glucose homeostasis via its role in the degradation of insulin, glucagon and IAPP (By similarity). Plays a role in the degradation and clearance of APP-derived amyloidogenic peptides that are secreted by neurons and microglia (PubMed:<a href="http://www.uniprot.org/citations/9830016" target="\_blank">>9830016</a>,  
PubMed:<a href="http://www.uniprot.org/citations/26394692"  
target="\_blank">>26394692</a>)  
(Probable). Degrades the natriuretic peptides ANP, BNP and CNP, inactivating their ability to raise intracellular cGMP (PubMed:<a href="http://www.uniprot.org/citations/21098034"  
target="\_blank">>21098034</a>). Also degrades an aberrant frameshifted 40-residue form of NPPA (fsNPPA) which is associated with familial atrial fibrillation in heterozygous patients (PubMed:<a href="http://www.uniprot.org/citations/21098034" target="\_blank">>21098034</a>). Involved in antigen processing. Produces both the N terminus and the C terminus of MAGEA3-derived antigenic peptide (EVDPIGHLY) that is presented to cytotoxic T lymphocytes by MHC class I.

#### Cellular Location

Cytoplasm, cytosol. Cell membrane {ECO:0000250|UniProtKB:P35559}. Secreted Note=Present at the cell surface of neuron cells. The membrane- associated isoform is approximately 5 kDa larger than the known cytosolic isoform

#### Tissue Location

Detected in brain and in cerebrospinal fluid (at protein level).



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with IDE antibody (Center)(Cat.#AP1455c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

#### IDE Antibody (Center) - Background

IDE belongs to a protease family responsible for intercellular peptide signalling. Though its role in the cellular processing of insulin has not yet been defined, insulin-degrading enzyme is thought to be involved in the termination of the insulin response.

#### IDE Antibody (Center) - References

- Vepsalainen,S.,J. Med. Genet. 44 (9), 606-608 (2007)  
Kim,M.,J. Biol. Chem. 282 (11), 7825-7832 (2007)  
Radulescu,R.T.,Int. J. Oncol. 30 (1), 73-80 (2007)  
Li,Q.,Cell 127 (2), 305-316 (2006)

#### IDE Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**IDE Antibody (Center) - Citations**

- [Promoting scientific standards in Germany.](#)
- [Complex formation between metabolic enzymes in tumor cells: unfolding the MDR1-IDE paradigm.](#)