

ZNF395 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14567c

Specification

ZNF395 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q9H8N7
Other Accession	NP_061130.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	54939
Antigen Region	293-322

ZNF395 Antibody (Center) - Additional Information

Gene ID 55893

Other Names

Zinc finger protein 395, HD-regulating factor 2, HDRF-2, Huntington disease gene regulatory region-binding protein 2, HD gene regulatory region-binding protein 2, HDBP-2, Papillomavirus regulatory factor 1, PRF-1, Papillomavirus-binding factor, ZNF395, HDBP2, PBF

Target/Specificity

This ZNF395 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 293-322 amino acids from the Central region of human ZNF395.

Dilution


WB~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



ZNF395 Antibody (Center) (Cat. #AP14567c) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the ZNF395 antibody detected the ZNF395 protein (arrow).

ZNF395 Antibody (Center) - Background

ZNF395 plays a role in papillomavirus genes transcription.

ZNF395 Antibody (Center) - References

- Tsukahara, T., et al. Cancer Sci. 100(1):47-53(2009)
- Sichtig, N., et al. Arch. Biochem. Biophys. 467(1):67-75(2007)
- Sichtig, N., et al. Arch. Biochem. Biophys. 464(1):90-99(2007)
- Tsukahara, T., et al. Cancer Res. 64(15):5442-5448(2004)
- Tanaka, K., et al. J. Biol. Chem. 279(8):7275-7286(2004)

cycles.

Precautions

ZNF395 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF395 Antibody (Center) - Protein Information

Name ZNF395

Synonyms HDBP2, PBF

Function

Plays a role in papillomavirus genes transcription.

Cellular Location

Cytoplasm. Nucleus. Note=May shuttle between nucleus and cytoplasm

Tissue Location

Widely expressed.

ZNF395 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)