

#### **CBX8 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP14665C

### **Specification**

#### **CBX8 Antibody (Center) - Product Information**

Application WB,E **Q9HC52** Primary Accession Other Accession NP 065700.1 Reactivity Mouse Host Rabbit Clonality **Polyclonal** Isotype Rabbit Ig Calculated MW 43396 Antigen Region 161-189

**CBX8 Antibody (Center) - Additional Information** 

#### **Gene ID 57332**

#### **Other Names**

Chromobox protein homolog 8, Polycomb 3 homolog, Pc3, hPc3, Rectachrome 1, CBX8, PC3, RC1

### **Target/Specificity**

This CBX8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 161-189 amino acids from the Central region of human CBX8.

## Dilution

WB~~1:1000

#### **Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

CBX8 Antibody (Center) is for research use only and not for use in diagnostic or



CBX8 Antibody (Center) (Cat. #AP14665c) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the CBX8 antibody detected the CBX8 protein (arrow).

# CBX8 Antibody (Center) - Background

Component of the Polycomb group (PcG) multiprotein PRC1 complex, a complex required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.

# **CBX8 Antibody (Center) - References**

Dietrich, N., et al. EMBO J. 26(6):1637-1648(2007)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Stelzl, U., et al. Cell 122(6):957-968(2005)
Obuse, C., et al. Genes Cells
9(2):105-120(2004)
Levine, S.S., et al. Mol. Cell. Biol. 22(17):6070-6078(2002)



therapeutic procedures.

**CBX8 Antibody (Center) - Protein Information** 

Name CBX8

Synonyms PC3, RC1

#### **Function**

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.

**Cellular Location** Nucleus.

# **CBX8 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture