

**Mouse Akt1 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14703a**

**Specification**

**Mouse Akt1 Antibody (N-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">P31750</a>
Other Accession	<a href="#">P31749</a> , <a href="#">NP_033782.1</a> , <a href="#">XP_001479205.1</a>
Reactivity	Mouse
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	55707
Antigen Region	22-49

**Mouse Akt1 Antibody (N-term) - Additional Information**

**Gene ID 11651**

**Other Names**

RAC-alpha serine/threonine-protein kinase, AKT1 kinase, Protein kinase B, PKB, Protein kinase B alpha, PKB alpha, Proto-oncogene c-Akt, RAC-PK-alpha, Thymoma viral proto-oncogene, Akt1, Akt, Rac

**Target/Specificity**

This Mouse Akt1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 22-49 amino acids from the N-terminal region of mouse Akt1.


**Dilution**

WB~1:1000  
IHC-P~1:10~50


**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**



Mouse Akt1 Antibody (N-term) (Cat. #AP14703a) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the Akt1 antibody detected the Akt1 protein (arrow).



Mouse Akt1 Antibody (N-term) (AP14703a) immunohistochemistry analysis in formalin fixed and paraffin embedded mouse gallbladder tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Mouse Akt1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Precautions

Mouse Akt1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### Mouse Akt1 Antibody (N-term) - Protein Information

**Name** Akt1

**Synonyms** Akt, Rac

### Function

AKT1 is one of 3 closely related serine/threonine-protein kinases (AKT1, AKT2 and AKT3) called the AKT kinase, and which regulate many processes including metabolism, proliferation, cell survival, growth and angiogenesis. This is mediated through serine and/or threonine phosphorylation of a range of downstream substrates. Over 100 substrate candidates have been reported so far, but for most of them, no isoform specificity has been reported (PubMed:<a href="http://www.uniprot.org/citations/11882383" target="\_blank">11882383</a>, PubMed:<a href="http://www.uniprot.org/citations/21620960" target="\_blank">21620960</a>, PubMed:<a href="http://www.uniprot.org/citations/21432781" target="\_blank">21432781</a>). AKT is responsible of the regulation of glucose uptake by mediating insulin-induced translocation of the SLC2A4/GLUT4 glucose transporter to the cell surface (PubMed:<a href="http://www.uniprot.org/citations/9415393" target="\_blank">9415393</a>). Phosphorylation of PTPN1 at 'Ser-50' negatively modulates its phosphatase activity preventing dephosphorylation of the insulin receptor and the attenuation of insulin signaling (PubMed:<a href="http://www.uniprot.org/citations/11579209" target="\_blank">11579209</a>). Phosphorylation of TBC1D4 triggers the binding of this effector to inhibitory 14-3-3 proteins, which is required for insulin-stimulated glucose transport (PubMed:<a href="http://www.uniprot.org/c

### Mouse Akt1 Antibody (N-term) - Background

This gene encodes the founding member of the Akt serine-threonine protein kinase gene family that also includes Akt2 and Akt3. This kinase is a major downstream effector of the phosphatidylinositol 3-kinase (PI3K) pathway that mediates the effects of various growth factors such as platelet-derived growth factor (PDGF), epidermal growth factor (EGF), insulin and insulin-like growth factor I (IGF-I). It is activated through recruitment to cellular membranes by PI3K lipid products and by phosphorylation by 3-phosphoinositide dependent kinase-1. It then further phosphorylates different downstream proteins in response to various extracellular signals and thus plays a pivotal role in mediating a variety of cellular processes, such as glucose metabolism, glycogen biosynthesis, protein synthesis and turn over, inflammatory response, cell survival (anti-apoptosis) and development. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq].

### Mouse Akt1 Antibody (N-term) - References

Chang, Z., et al. Dev. Biol. 347(2):384-391(2010)  
Zhou, F., et al. Am. J. Pathol. 177(4):2124-2133(2010)  
Sun, X., et al. Proc. Natl. Acad. Sci. U.S.A. 107(39):16887-16892(2010)  
Goncalves, M.D., et al. PLoS ONE 5 (9), E12707 (2010) :  
Bremer, J., et al. PLoS ONE 5 (8), E12450 (2010) :

itations/11994271" target="\_blank">11994271

AKT regulates also the storage of glucose in the form of glycogen by phosphorylating GSK3A at 'Ser-21' and GSK3B at 'Ser-9', resulting in inhibition of its kinase activity (PubMed: [22057101](http://www.uniprot.org/citations/22057101)). Phosphorylation of GSK3 isoforms by AKT is also thought to be one mechanism by which cell proliferation is driven (PubMed: [22057101](http://www.uniprot.org/citations/22057101)). AKT regulates also cell survival via the phosphorylation of MAP3K5 (apoptosis signal-related kinase). Phosphorylation of 'Ser-83' decreases MAP3K5 kinase activity stimulated by oxidative stress and thereby prevents apoptosis. AKT mediates insulin-stimulated protein synthesis by phosphorylating TSC2 at 'Ser- 939' and 'Thr-1462', thereby activating mTORC1 signaling and leading to both phosphorylation of 4E-BP1 and in activation of RPS6KB1. AKT is involved in the phosphorylation of members of the FOXO factors (Forkhead family of transcription factors), leading to binding of 14-3-3 proteins and cytoplasmic localization. In particular, FOXO1 is phosphorylated at 'Thr-24', 'Ser-256' and 'Ser-319'. FOXO3 and FOXO4 are phosphorylated on equivalent sites. AKT has an important role in the regulation of NF-kappa-B-dependent gene transcription and positively regulates the activity of CREB1 (cyclic AMP (cAMP)-response element binding protein). The phosphorylation of CREB1 induces the binding of accessory proteins that are necessary for the transcription of pro-survival genes such as BCL2 and MCL1 (By similarity). AKT phosphorylates 'Ser-454' on ATP citrate lyase (ACLY), thereby potentially regulating ACLY activity and fatty acid synthesis (By similarity). Activates the 3B isoform of cyclic nucleotide phosphodiesterase (PDE3B) via phosphorylation of 'Ser-273', resulting in reduced cyclic AMP levels and inhibition of lipolysis (PubMed: [10454575](http://www.uniprot.org/citations/10454575)). Phosphorylates PIKFYVE on 'Ser-318', which results in increased PI(3)P-5 activity (By similarity). The Rho GTPase- activating protein DLC1 is another substrate and its phosphorylation is implicated in the

regulation cell proliferation and cell growth (By similarity). AKT plays a role as key modulator of the AKT-mTOR signaling pathway controlling the tempo of the process of newborn neurons integration during adult neurogenesis, including correct neuron positioning, dendritic development and synapse formation (PubMed:<a href="http://www.uniprot.org/citations/19778506" target="\_blank">19778506</a>). Signals downstream of phosphatidylinositol 3-kinase (PI(3)K) to mediate the effects of various growth factors such as platelet-derived growth factor (PDGF), epidermal growth factor (EGF), insulin and insulin-like growth factor I (IGF-I) (PubMed:<a href="http://www.uniprot.org/citations/11282895" target="\_blank">11282895</a>, PubMed:<a href="http://www.uniprot.org/citations/18288188" target="\_blank">18288188</a>). AKT mediates the antiapoptotic effects of IGF-I (PubMed:<a href="http://www.uniprot.org/citations/11282895" target="\_blank">11282895</a>). Essential for the SPATA13-mediated regulation of cell migration and adhesion assembly and disassembly (By similarity). May be involved in the regulation of the placental development (PubMed:<a href="http://www.uniprot.org/citations/12783884" target="\_blank">12783884</a>). Phosphorylates STK4/MST1 at 'Thr-120' and 'Thr-387' leading to inhibition of its kinase activity, nuclear translocation, autophosphorylation and ability to phosphorylate FOXO3. Phosphorylates STK3/MST2 at 'Thr-117' and 'Thr-384' leading to inhibition of its cleavage, kinase activity, autophosphorylation at Thr-180, binding to RASSF1 and nuclear translocation. Phosphorylates SRPK2 and enhances its kinase activity towards SRSF2 and ACIN1 and promotes its nuclear translocation. Phosphorylates RAF1 at 'Ser-259' and negatively regulates its activity. Phosphorylation of BAD stimulates its pro-apoptotic activity. Phosphorylates KAT6A at 'Thr-369' and this phosphorylation inhibits the interaction of KAT6A with PML and negatively regulates its acetylation activity towards p53/TP53. Phosphorylates palladin (PALLD), modulating cytoskeletal organization and cell motility. Phosphorylates prohibitin (PHB), playing an important role in cell metabolism and

proliferation. Phosphorylates CDKN1A, for which phosphorylation at 'Thr-145' induces its release from CDK2 and cytoplasmic relocalization. These recent findings indicate that the AKT1 isoform has a more specific role in cell motility and proliferation. Phosphorylates CLK2 thereby controlling cell survival to ionizing radiation (By similarity). Phosphorylates PCK1 at 'Ser-90', reducing the binding affinity of PCK1 to oxaloacetate and changing PCK1 into an atypical protein kinase activity using GTP as donor (By similarity). Also acts as an activator of TMEM175 potassium channel activity in response to growth factors: forms the lysoK(GF) complex together with TMEM175 and acts by promoting TMEM175 channel activation, independently of its protein kinase activity (PubMed:<a href="http://www.uniprot.org/citations/32228865" target="\_blank">32228865</a>).

#### **Cellular Location**

Cytoplasm. Nucleus. Cell membrane. Note=Nucleus after activation by integrin-linked protein kinase 1 (ILK1) (By similarity). Nuclear translocation is enhanced by interaction with TCL1A. Phosphorylation on Tyr-176 by TNK2 results in its localization to the cell membrane where it is targeted for further phosphorylations on Thr-308 and Ser-473 leading to its activation and the activated form translocates to the nucleus. Colocalizes with WDFY2 in intracellular vesicles {ECO:0000250|UniProtKB:P31749}

#### **Tissue Location**

Widely expressed. Low levels found in liver with slightly higher levels present in thymus and testis

#### **Mouse Akt1 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)