

**GPC3 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14822b**

**Specification**

**GPC3 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P51654</a>
Other Accession	<a href="#">NP_004475.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	65563
Antigen Region	465-494

**GPC3 Antibody (C-term) - Additional Information**

**Gene ID** 2719

**Other Names**

Glypican-3, GTR2-2, Intestinal protein OCI-5, MXR7, Secreted glypican-3, GPC3, OCI5

**Target/Specificity**

This GPC3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 465-494 amino acids from the C-terminal region of human GPC3.

**Dilution**

WB~~1:1000

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GPC3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



GPC3 Antibody (C-term) (Cat. #AP14822b) western blot analysis in NCI-H292 cell line lysates (35ug/lane). This demonstrates the GPC3 antibody detected the GPC3 protein (arrow).

**GPC3 Antibody (C-term) - Background**

Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi-Behmel syndrome, also known as Simpson dysmorphia syndrome. Alternative splicing results in

**GPC3 Antibody (C-term) - Protein Information****Name** GPC3**Synonyms** OCI5**Function**

Cell surface proteoglycan that bears heparan sulfate (PubMed:<a href="http://www.uniprot.org/citations/14610063" target="\_blank">14610063</a>). Negatively regulates the hedgehog signaling pathway when attached via the GPI-anchor to the cell surface by competing with the hedgehog receptor PTC1 for binding to hedgehog proteins (By similarity). Binding to the hedgehog protein SHH triggers internalization of the complex by endocytosis and its subsequent lysosomal degradation (By similarity). Positively regulates the canonical Wnt signaling pathway by binding to the Wnt receptor Frizzled and stimulating the binding of the Frizzled receptor to Wnt ligands (PubMed:<a href="http://www.uniprot.org/citations/16227623" target="\_blank">16227623</a>, PubMed:<a href="http://www.uniprot.org/citations/24496449" target="\_blank">24496449</a>). Positively regulates the non- canonical Wnt signaling pathway (By similarity). Binds to CD81 which decreases the availability of free CD81 for binding to the transcriptional repressor HHEX, resulting in nuclear translocation of HHEX and transcriptional repression (By similarity). Inhibits the dipeptidyl peptidase activity of DPP4 (PubMed:<a href="http://www.uniprot.org/citations/17549790" target="\_blank">17549790</a>). Plays a role in limb patterning and skeletal development by controlling the cellular response to BMP4 (By similarity). Modulates the effects of growth factors BMP2, BMP7 and FGF7 on renal branching morphogenesis (By similarity). Required for coronary vascular development (By similarity). Plays a role in regulating cell movements during gastrulation (By similarity).

**Cellular Location**

Cell membrane; Lipid-anchor, GPI-anchor {ECO:0000250|UniProtKB:P13265}; Extracellular side

multiple transcript variants.

**GPC3 Antibody (C-term) - References**

Lai, J.P., et al. Hepatology 52(5):1680-1689(2010)  
Sakurai, M., et al. Gynecol. Oncol. 119(2):332-336(2010)  
Zynger, D.L., et al. Histopathology 56(6):750-757(2010)  
Davoodi, J., et al. Proteomics 7(13):2300-2310(2007)  
Hsu, H.C., et al. Cancer Res. 57(22):5179-5184(1997)

{ECO:0000250|UniProtKB:P13265}

**Tissue Location**

Highly expressed in lung, liver and kidney.

**GPC3 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)