

**CDH10 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1482b**

**Specification**

**CDH10 Antibody (C-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">Q9Y6N8</a>
Other Accession	<a href="#">P70408</a> , <a href="#">P79995</a>
Reactivity	Human, Mouse, Rat
Predicted Host	Chicken
Clonality	Rabbit
Isotype	Polyclonal
Antigen Region	Rabbit Ig
	495-523

**CDH10 Antibody (C-term) - Additional Information**

**Gene ID** 1008

**Other Names**

Cadherin-10, T2-cadherin, CDH10

**Target/Specificity**

This CDH10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 495-523 amino acids from the C-terminal region of human CDH10.

**Dilution**

WB~~1:1000  
IHC-P~~1:10~50  
FC~~1:10~50

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CDH10 Antibody (C-term) is for research



All lanes : Anti-CDH10 Antibody (C-term) at 1:2000 dilution Lane 1: SK-BR-3 whole cell lysate Lane 2: NCI-H460 whole cell lysate Lane 3: F9 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 88 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-CDH10 Antibody (C-term) at 1:1000 dilution + C6 whole cell lysate

use only and not for use in diagnostic or therapeutic procedures.

#### CDH10 Antibody (C-term) - Protein Information

**Name** CDH10

#### Function

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types.

#### Cellular Location

Cell membrane; Single-pass type I membrane protein

#### Tissue Location

Predominantly expressed in brain. Also found in adult and fetal kidney. Very low levels detected in prostate and fetal lung.

#### CDH10 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L),  
Peroxidase conjugated at 1/10000 dilution.  
Predicted band size : 88 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human brain tissue reacted with CDH10 antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of NCI-H460 cells using Cadherin 10 (CDH10) Antibody (C-term) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### CDH10 Antibody (C-term) - Background

CDH10 is a type II classical cadherin from the cadherin superfamily, integral membrane proteins that mediate calcium-dependent cell-cell adhesion. Mature cadherin proteins are composed of a large N-terminal extracellular

domain, a single membrane-spanning domain, and a small, highly conserved C-terminal cytoplasmic domain. The extracellular domain consists of 5 subdomains, each containing a cadherin motif, and appears to determine the specificity of the protein's homophilic cell adhesion activity. Type II (atypical) cadherins are defined based on their lack of a HAV cell adhesion recognition sequence specific to type I cadherins. This particular cadherin is predominantly expressed in brain and is putatively involved in synaptic adhesions, axon outgrowth and guidance.

#### **CDH10 Antibody (C-term) - References**

Kools,P., FEBS Lett. 452 (3), 328-334 (1999)

#### **CDH10 Antibody (C-term) - Citations**

- [Alterations of type II classical cadherin Cadherin-10 \(CDH10\) is associated with pancreatic ductal adenocarcinomas.](#)