

ECSIT Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14858b

Specification

ECSIT Antibody (C-term) - Product Information

Application	IF, WB, E
Primary Accession	Q9BO95
Other Accession	NP_001135936.1 , NP_057665.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	396-425

ECSIT Antibody (C-term) - Additional Information

Gene ID 51295

Other Names

Evolutionarily conserved signaling intermediate in Toll pathway, mitochondrial, Protein SITPEC, ECSIT

Target/Specificity

This ECSIT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 396-425 amino acids from the C-terminal region of human ECSIT.

Dilution

IF~~1:10~50

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ECSIT Antibody (C-term) is for research use



Fluorescent confocal image of HeLa cell stained with ECSIT Antibody (C-term)(Cat#AP14858b). HeLa cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with ECSIT primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). ECSIT immunoreactivity is localized to Cytoplasm and Nucleus significantly.

only and not for use in diagnostic or therapeutic procedures.

ECSIT Antibody (C-term) - Protein Information

Name ECSIT

Function

Adapter protein of the Toll-like and IL-1 receptor signaling pathway that is involved in the activation of NF-kappa-B via MAP3K1. Promotes proteolytic activation of MAP3K1. Involved in the BMP signaling pathway. Required for normal embryonic development (By similarity).


Cellular Location

Cytoplasm. Nucleus. Mitochondrion

ECSIT Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



ECSIT Antibody (C-term) (Cat. #AP14858b) western blot analysis in Jurkat,ZR-75-1 cell line lysates (35ug/lane).This demonstrates the ECSIT antibody detected the ECSIT protein (arrow).

ECSIT Antibody (C-term) - Background

Adapter protein of the Toll-like and IL-1 receptor signaling pathway that is involved in the activation of NF-kappa-B via MAP3K1. Promotes proteolytic activation of MAP3K1. Involved in the BMP signaling pathway. Required for normal embryonic development (By similarity). Required for efficient assembly of mitochondrial NADH:ubiquinone oxidoreductase.

ECSIT Antibody (C-term) - References

Vogel, R.O., et al. Genes Dev. 21(5):615-624(2007)
Xiao, C., et al. Genes Dev. 17(23):2933-2949(2003)
Kopp, E., et al. Genes Dev. 13(16):2059-2071(1999)