

**TLR2 Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1502A**

### Specification

#### TLR2 Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">O60603</a>
Other Accession	<a href="#">Q95M53</a>
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	212-242

#### TLR2 Antibody (N-term) - Additional Information

##### Gene ID 7097

##### Other Names

Toll-like receptor 2, Toll/interleukin-1 receptor-like protein 4, CD282, TLR2, TIL4

##### Target/Specificity

This TLR2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 212-242 amino acids from the N-terminal region of human TLR2.

##### Dilution

WB~~1:1000  
IHC-P~~1:10~50  
FC~~1:10~50

##### Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

##### Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

TLR2 Antibody (N-term) is for research use



Anti-TLR2 Antibody (C226) at 1:1000 dilution + Ramos whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 90 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with TLR2 antibody (N-term) (Cat.#AP1502a), which was peroxidase-conjugated to the secondary antibody, followed by DAB

only and not for use in diagnostic or therapeutic procedures.

#### TLR2 Antibody (N-term) - Protein Information


**Name** TLR2

**Synonyms** TIL4

#### Function

Cooperates with LY96 to mediate the innate immune response to bacterial lipoproteins and other microbial cell wall components. Cooperates with TLR1 or TLR6 to mediate the innate immune response to bacterial lipoproteins or lipopeptides (PubMed:<a href="http://www.uniprot.org/citations/21078852" target="\_blank">21078852</a>, PubMed:<a href="http://www.uniprot.org/citations/17889651" target="\_blank">17889651</a>). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. May also activate immune cells and promote apoptosis in response to the lipid moiety of lipoproteins (PubMed:<a href="http://www.uniprot.org/citations/10426995" target="\_blank">10426995</a>, PubMed:<a href="http://www.uniprot.org/citations/10426996" target="\_blank">10426996</a>). Recognizes mycoplasmal macrophage-activating lipopeptide-2kD (MALP-2), soluble tuberculosis factor (STF), phenol-soluble modulin (PSM) and *B.burgdorferi* outer surface protein A lipoprotein (OspA-L) cooperatively with TLR6 (PubMed:<a href="http://www.uniprot.org/citations/11441107" target="\_blank">11441107</a>). Stimulation of monocytes in vitro with *M.tuberculosis* PstS1 induces p38 MAPK and ERK1/2 activation primarily via this receptor, but also partially via TLR4 (PubMed:<a href="http://www.uniprot.org/citations/16622205" target="\_blank">16622205</a>). MAPK activation in response to bacterial peptidoglycan also occurs via this receptor (PubMed:<a href="http://www.uniprot.org/citations/16622205" target="\_blank">16622205</a>). Acts as a receptor for *M.tuberculosis* lipoproteins LprA, LprG, LpqH and PstS1, some lipoproteins are dependent on other

staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of CEM cells using TLR2 Antibody (N-term) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### TLR2 Antibody (N-term) - Background

The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from *Drosophila* to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is expressed most abundantly in peripheral blood leukocytes, and mediates host response to Gram-positive bacteria and yeast via stimulation of NF-kappaB.

#### TLR2 Antibody (N-term) - References

Meng, G., et al., *J. Biol. Chem.* 278(41):39822-39829 (2003). Sandor, F., et al., *J. Cell Biol.* 162(6):1099-1110 (2003). Wang, X., et al., *FASEB J.* 17(12):1727-1729 (2003). Huang, L.Y., et al., *J. Immunol.*

coreceptors (TLR1, CD14 and/or CD36); the lipoproteins act as agonists to modulate antigen presenting cell functions in response to the pathogen (PubMed:<a href ="<http://www.uniprot.org/citations/19362712>" target="\_blank">19362712</a>). M.tuberculosis HSP70 (dnaK) but not HSP65 (groEL-2) acts via this protein to stimulate NF-kappa-B expression (PubMed:<a href="<http://www.uniprot.org/citations/15809303>" target="\_blank">15809303</a>). Recognizes M.tuberculosis major T-antigen EsxA (ESAT-6) which inhibits downstream MYD88-dependent signaling (shown in mouse) (By similarity). Forms activation clusters composed of several receptors depending on the ligand, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway. Forms the cluster TLR2:TLR6:CD14:CD36 in response to diacylated lipopeptides and TLR2:TLR1:CD14 in response to triacylated lipopeptides (PubMed:<a href="http://www.uniprot.org/citations/16880211" target="\_blank">16880211</a>). Required for normal uptake of M.tuberculosis, a process that is inhibited by M.tuberculosis LppM (By similarity).

### Cellular Location

#### Membrane

{ECO:0000250|UniProtKB:Q9QUN7}; Single-pass type I membrane protein. Cytoplasmic vesicle, phagosome membrane  
{ECO:0000250|UniProtKB:Q9QUN7};  
Single-pass type I membrane protein.  
Membrane raft. Note=Does not reside in lipid rafts before stimulation but accumulates increasingly in the raft upon the presence of the microbial ligand. In response to diacylated lipoproteins, TLR2:TLR6 heterodimers are recruited in lipid rafts, this recruitment determines the intracellular targeting to the Golgi apparatus. Triacylated lipoproteins induce the same mechanism for TLR2:TLR1 heterodimers.

### Tissue Location

Highly expressed in peripheral blood leukocytes, in particular in monocytes, in bone marrow, lymph node and in spleen. Also detected in lung and in fetal liver. Levels are low in other tissues

171(3):1441-1446 (2003). Takeuchi, J., et al., Biochem. Biophys. Res. Commun. 306(3):674-679 (2003).

## TLR2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## TLR2 Antibody (N-term) - Citations

- [TLR2 antagonism attenuates the hippocampal neuronal damage in a murine model of sleep apnea via inhibiting neuroinflammation and oxidative stress](#)
- [Migration ability and Toll-like receptor expression of human mesenchymal stem cells improves significantly after three-dimensional culture.](#)
- [Bacterial peptidoglycan breaks down intestinal tolerance via mast cell activation: the role of TLR2 and NOD2.](#)