

M TLR4 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1504a

Specification

M TLR4 Antibody (N-term) - Product Information

Application	WB, E
Primary Accession	Q9QUK6
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	25-55

M TLR4 Antibody (N-term) - Additional Information

Gene ID 21898

Other Names

Toll-like receptor 4, CD284, Tlr4, Lps

Target/Specificity

This Mouse TLR4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 25-55 amino acids from the N-terminal region of mouse TLR4.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

M TLR4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of anti-mTLR4 Pab (Cat. #AP1504a) in mouse spleen cell lysate. mTLR4 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

M TLR4 Antibody (N-term) - Background

TLR4, a type I membrane protein that belongs to the Toll-like receptor family, cooperates with LY96 and CD14 to mediate the innate immune response to bacterial lipopolysaccharide (LPS). It acts via MyD88, TIRAP and TRAF6, leading to NF- κ B activation, cytokine secretion and the inflammatory response. TLR4 Belongs to the lipopolysaccharide (LPS) receptor, a multi-protein complex containing at least CD14, LY96 and TLR. TLR4 binds to LY96 via the extracellular domain, and to MyD88 and TIRAP via their respective TIR domains. The protein contains 19 leucine-rich (LRR) repeats, and it is highly expressed in heart, spleen, lung and muscle. Lower levels are found in liver and kidney. Interstrain analyses reveal that TLR4 is a polymorphic protein and that the extracellular domain is far more variable than the cytoplasmic domain, which is variable at the C-terminal.

M TLR4 Antibody (N-term) - Protein Information**Name** Tlr4**Synonyms** Lps**Function**

Cooperates with LY96 and CD14 to mediate the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed:9851930, PubMed:9989976, PubMed:20133493). Acts via MYD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:24380872). Also involved in LPS- independent inflammatory responses triggered by free fatty acids, such as palmitate. In complex with TLR6, promotes sterile inflammation in monocytes/macrophages in response to oxidized low-density lipoprotein (oxLDL) or amyloid-beta 42. In this context, the initial signal is provided by oxLDL- or amyloid-beta 42-binding to CD36. This event induces the formation of a heterodimer of TLR4 and TLR6, which is rapidly internalized and triggers inflammatory response, leading to the NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway, as well as IL1B secretion. Binds electronegative LDL (LDL-) and mediates the cytokine release induced by LDL- (By similarity). Activated by the signaling pathway regulator NMI which acts as damage- associated molecular patterns (DAMPs) in response to cell injury or pathogen invasion, therefore promoting nuclear factor NF-kappa-B activation (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Early endosome {ECO:0000250|UniProtKB:O00206}. Cell projection, ruffle. Note=Upon complex formation with CD36 and TLR6, internalized

through dynamin-dependent endocytosis
Colocalizes with RFTN1 at cell membrane
and then together with RFTN1 moves to
endosomes, upon lipopolysaccharide
stimulation
{ECO:0000250|UniProtKB:O00206}

Tissue Location

Expressed in macrophages (at protein level)
(PubMed:28098138). Highly expressed in
heart, spleen, lung and muscle Lower levels
are found in liver and kidney
(PubMed:23812099)

M TLR4 Antibody (N-term) - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

M TLR4 Antibody (N-term) - Citations

- [Moderate prenatal alcohol exposure suppresses the TLR4-mediated innate immune response in the hippocampus of young rats.](#)
- [Inhibition of MyD88 Signaling Skews Microglia/Macrophage Polarization and Attenuates Neuronal Apoptosis in the Hippocampus After Status Epilepticus in Mice.](#)
- [Pretreatment of Huaiqihuang extractum protects against cisplatin-induced nephrotoxicity.](#)
- [Resveratrol alleviates lysophosphatidylcholine-induced damage and inflammation in vascular endothelial cells.](#)
- [Atherogenic high cholesterol/high fat diet induces TLRs-associated pulmonary inflammation in C57BL/6J mice.](#)
- [Lead exposure induced microgliosis and astrogliosis in hippocampus of young mice potentially by triggering TLR4-MyD88-NF \$\kappa\$ B signaling cascades.](#)
- [Efficacy of atorvastatin on hippocampal neuronal damage caused by chronic intermittent hypoxia: involving TLR4 and its downstream signaling pathway.](#)
- [Uncontrolled inflammation induced by AEG-1 promotes gastric cancer and poor prognosis.](#)
- [Caspase-8 promotes NLRP1/NLRP3 inflammasome activation and IL-1 \$\beta\$ production in acute glaucoma.](#)
- [The tumor suppressor p15Ink4b regulates the differentiation and maturation of conventional dendritic cells.](#)
- [Arteriogenesis requires toll-like receptor 2 and 4 expression in bone-marrow derived cells.](#)