

NEIL2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16044a

Specification

NEIL2 Antibody (N-term) - Product Information

Application	WB, E
Primary Accession	Q969S2
Other Accession	NP_001129220.1 , NP_001129218.1 , NP_659480.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	36826
Antigen Region	67-96

NEIL2 Antibody (N-term) - Additional Information

Gene ID 252969

Other Names

Endonuclease 8-like 2, 322-, DNA glycosylase/AP lyase Neil2, DNA-(apurinic or apyrimidinic site) lyase Neil2, Endonuclease VIII-like 2, Nei homolog 2, NEH2, Nei-like protein 2, NEIL2

Target/Specificity

This NEIL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 67-96 amino acids from the N-terminal region of human NEIL2.

Dilution


WB~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



NEIL2 Antibody (N-term) (Cat. #AP16044a) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the NEIL2 antibody detected the NEIL2 protein (arrow).

NEIL2 Antibody (N-term) - Background

NEIL2 belongs to a class of DNA glycosylases homologous to the bacterial Fpg/Nei family. These glycosylases initiate the first step in base excision repair by cleaving bases damaged by reactive oxygen species and introducing a DNA strand break via the associated lyase reaction (Bandaru et al., 2002 [PubMed 12509226])

NEIL2 Antibody (N-term) - References

Briggs, F.B., et al. Am. J. Epidemiol. 172(2):217-224(2010)
Monsees, G.M., et al. Breast Cancer Res. Treat. (2010) In press :
Grin, I.R., et al. Biochem. Biophys. Res. Commun. 394(1):100-105(2010)
Kinslow, C.J., et al. Genes Chromosomes Cancer 47(11):923-932(2008)
Zhai, X., et al. Clin. Cancer Res.

Precautions

NEIL2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

14(13):4345-4352(2008)

NEIL2 Antibody (N-term) - Protein Information**Name** NEIL2**Function**

Involved in base excision repair of DNA damaged by oxidation or by mutagenic agents. Has DNA glycosylase activity towards 5-hydroxyuracil and other oxidized derivatives of cytosine with a preference for mismatched double-stranded DNA (DNA bubbles). Has low or no DNA glycosylase activity towards thymine glycol, 2-hydroxyadenine, hypoxanthine and 8-oxoguanine. Has AP (apurinic/apyrimidinic) lyase activity and introduces nicks in the DNA strand. Cleaves the DNA backbone by beta-delta elimination to generate a single-strand break at the site of the removed base with both 3'- and 5'-phosphates.

Cellular Location

Nucleus.

Tissue Location

Detected in testis, skeletal muscle, heart, brain, placenta, lung, pancreas, kidney and liver

NEIL2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)