

GATA3 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16085c

Specification

GATA3 Antibody (Center) - Product Information

Application	WB, E
Primary Accession	P23771
Other Accession	P23772 , P23825 , NP_002042.1
Reactivity	Human, Mouse
Predicted	Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	47916
Antigen Region	235-263

GATA3 Antibody (Center) - Additional Information

Gene ID 2625

Other Names

Trans-acting T-cell-specific transcription factor GATA-3, GATA-binding factor 3, GATA3

Target/Specificity

This GATA3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 235-263 amino acids from the Central region of human GATA3.

Dilution

WB~1:1000


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



GATA3 Antibody (Center) (Cat. #AP16085c) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the GATA3 antibody detected the GATA3 protein (arrow).



Western blot analysis of GATA3 (arrow) using rabbit polyclonal GATA3 Antibody (Center) (Cat. #AP16085c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the GATA3 gene.

GATA3 Antibody (Center) - Background

This gene encodes a protein which belongs to the GATA

GATA3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

GATA3 Antibody (Center) - Protein Information

Name GATA3

Function

Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper 2 (Th2) differentiation process following immune and inflammatory responses.

Cellular Location

Nucleus.

Tissue Location

T-cells and endothelial cells.

family of transcription factors. The protein contains two GATA-type zinc fingers and is an important regulator of T-cell development and plays an important role in endothelial cell biology. Defects in this gene are the cause of hypoparathyroidism with sensorineural deafness and renal dysplasia.

GATA3 Antibody (Center) - References

Cook, K.D., et al. J. Immunol. 185(6):3209-3216(2010)
Schuurhof, A., et al. Pediatr. Pulmonol. 45(6):608-613(2010)
Kemp, K.L., et al. J. Immunol. 184(8):4178-4184(2010)
Davila, S., et al. Genes Immun. 11(3):232-238(2010)
Yao, X., et al. Chin. Med. J. 123(6):739-744(2010)

GATA3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)