

CLC Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16125b

Specification

CLC Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q05315
Other Accession	NP_001819.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	16453
Antigen Region	91-119

CLC Antibody (C-term) - Additional Information

Gene ID 1178

Other Names

Galectin-10, Gal-10, Charcot-Leyden crystal protein, CLC, Eosinophil lysophospholipase, Lysolecithin acylhydrolase, CLC, LGALS10, LGALS10A

Target/Specificity

This CLC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 91-119 amino acids from the C-terminal region of human CLC.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CLC Antibody (C-term) is for research use only and not for use in diagnostic or



Western blot analysis of CLC (arrow) using rabbit polyclonal CLC Antibody (C-term) (Cat. #AP16125b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CLC gene.

CLC Antibody (C-term) - Background

Lysophospholipases are enzymes that act on biological membranes to regulate the multifunctional lysophospholipids. The protein encoded by this gene is a lysophospholipase expressed in eosinophils and basophils. It hydrolyzes lysophosphatidylcholine to glycerophosphocholine and a free fatty acid. This protein may possess carbohydrate or IgE-binding activities. It is both structurally and functionally related to the galectin family of beta-galactoside binding proteins. It may be associated with inflammation and some myeloid leukemias.

CLC Antibody (C-term) - References

Davila, S., et al. Genes Immun. 11(3):232-238(2010)
Bryborn, M., et al. Allergy 65(2):220-228(2010)

therapeutic procedures.

CLC Antibody (C-term) - Protein Information

Name CLC

Synonyms LGALS10, LGALS10A

Function

Regulates immune responses through the recognition of cell- surface glycans. Essential for the anergy and suppressive function of CD25-positive regulatory T-cells (Treg).

Cellular Location

Cytoplasm, cytosol. Cytoplasmic granule. Note=Localized in granules from where it may be secreted or transported to other locations in the cell

Tissue Location

Expressed abundantly in the bone marrow. Expressed exclusively by eosinophils and basophils. Not detected in monocytes and neutrophils. Expressed in CD25-positive regulatory T-cells (Treg) (at protein level). Found in intestinal tissue from patients with Celiac disease, expression is directly related to the histological grade of mucosal damage and to the number of eosinophils found in the duodenal lesion (at protein level). Found in sputum of patients with eosinophilic inflammatory diseases such as asthma (at protein level)

De Re, V., et al. Ann. N. Y. Acad. Sci. 1173, 357-364 (2009) :
Than, N.G., et al. Proc. Natl. Acad. Sci. U.S.A. 106(24):9731-9736(2009)
Kubach, J., et al. Blood 110(5):1550-1558(2007)

CLC Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)