

CEP72 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16441c

Specification

CEP72 Antibody (Center) - Product Information

Application	WB, E
Primary Accession	Q9P209
Other Accession	NP_060610.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	71718
Antigen Region	231-259

CEP72 Antibody (Center) - Additional Information

Gene ID 55722

Other Names

Centrosomal protein of 72 kDa, Cep72, CEP72, KIAA1519

Target/Specificity

This CEP72 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 231-259 amino acids from the Central region of human CEP72.

Dilution

WB~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CEP72 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.



CEP72 Antibody (Center) (Cat. #AP16441c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the CEP72 antibody detected the CEP72 protein (arrow).

CEP72 Antibody (Center) - Background

CEP72 is a member of the leucine-rich-repeat (LRR) superfamily of proteins. The protein is localized to the centrosome, a non-membraneous organelle that functions as the major microtubule-organizing center in animal cells.

CEP72 Antibody (Center) - References

McGovern, D.P., et al. *Nat. Genet.* 42(4):332-337(2010)
Briggs, F.B., et al. *Genes Immun.* 11(3):199-208(2010)
Oshimori, N., et al. *EMBO J.* 28(14):2066-2076(2009)
Xin, X., et al. *Genome Res.* 19(7):1262-1269(2009)
Olsen, J.V., et al. *Cell* 127(3):635-648(2006)

CEP72 Antibody (Center) - Protein Information**Name** CEP72**Synonyms** KIAA1519**Function**

Involved in the recruitment of key centrosomal proteins to the centrosome. Provides centrosomal microtubule-nucleation activity on the gamma-tubulin ring complexes (gamma-TuRCs) and has critical roles in forming a focused bipolar spindle, which is needed for proper tension generation between sister chromatids. Required for localization of KIZ, AKAP9 and gamma-tubulin ring complexes (gamma-TuRCs) (PubMed:19536135). Involved in centriole duplication. Required for CDK5RAP22, CEP152, WDR62 and CEP63 centrosomal localization and promotes the centrosomal localization of CDK2 (PubMed:26297806).

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite. Note=Localizes to the centrosome and centrosome-surrounding particles throughout the cell cycle. These particles disappear after microtubules are depolymerized using nocodazole, suggesting that CEP72-associating particles localize in a microtubule- dependent manner

CEP72 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)