

RAB41 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16499b

Specification

RAB41 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q5JT25
Other Accession	NP_001027898.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	25038
Antigen Region	137-165

RAB41 Antibody (C-term) - Additional Information

Gene ID 347517

Other Names

Ras-related protein Rab-41, RAB41

Target/Specificity

This RAB41 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 137-165 amino acids from the C-terminal region of human RAB41.

Dilution

WB~~1:1000

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAB41 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



RAB41 Antibody (C-term) (Cat. #AP16499b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the RAB41 antibody detected the RAB41 protein (arrow).

RAB41 Antibody (C-term) - Background

This gene encodes a small GTP-binding protein that belongs to the largest family within the Ras superfamily. These proteins function as regulators of membrane trafficking. They cycle between inactive GDP-bound and activated GTP-bound states, which is controlled by GTP hydrolysis-activating proteins (GAPs). This family member can be activated by the GAP protein RN-Tre, and it is localized to the Golgi complex.

RAB41 Antibody (C-term) - References

Haas, A.K., et al. Nat. Cell Biol. 7(9):887-893(2005)
Pereira-Leal, J.B., et al. J. Mol. Biol. 313(4):889-901(2001)

RAB41 Antibody (C-term) - Protein Information

Name RAB41

Function

Required for normal Golgi ribbon organization and ER-to-Golgi trafficking.

Cellular Location

Cytoplasm. Note=punctate localization concentrated in ruffled regions at the cell periphery

Tissue Location

Widely expressed in brain, testis, lung, heart, ovary, colon, kidney, uterus and spleen but not in liver

RAB41 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)