

TGFBI Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16708a

Specification

TGFBI Antibody (N-term) - Product Information

Application	WB, FC, E
Primary Accession	Q15582
Other Accession	Q95215 , Q11780 , P82198 , NP_000349.1 , P55906
Reactivity	Human
Predicted	Bovine, Mouse, Pig, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	74681
Antigen Region	106-135

TGFBI Antibody (N-term) - Additional Information

Gene ID 7045

Other Names

Transforming growth factor-beta-induced protein ig-h3, Beta ig-h3, Kerato-epithelin, RGD-containing collagen-associated protein, RGD-CAP, TGFBI, BIGH3

Target/Specificity

This TGFBI antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 106-135 amino acids from the N-terminal region of human TGFBI.

Dilution


WB~~1:1000
 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2



Western blot analysis of TGFBI (arrow) using rabbit polyclonal TGFBI Antibody (N-term) (Cat. #AP16708a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the TGFBI gene.



TGFBI Antibody (N-term) (Cat. #AP16708a) flow cytometric analysis of U251 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TGFBI Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TGFBI Antibody (N-term) - Protein Information

Name TGFBI

Synonyms BIGH3

Function

Plays a role in cell adhesion (PubMed:8024701). May play a role in cell-collagen interactions (By similarity).

Cellular Location

Secreted. Secreted, extracellular space, extracellular matrix Note=May be associated both with microfibrils and with the cell surface (PubMed:8077289).

Tissue Location

Highly expressed in the corneal epithelium (PubMed:27609313, PubMed:8077289).

Expressed in heart, placenta, lung, liver, skeletal muscle, kidney and pancreas

(PubMed:8077289)

TGFBI Antibody (N-term) - Background

This gene encodes an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. This protein plays a role in cell-collagen interactions and may be involved in endochondrial bone formation in cartilage. The protein is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in this gene are associated with multiple types of corneal dystrophy.

TGFBI Antibody (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Edelstein, S.L., et al. Cornea 29(6):698-700(2010)
Romero, P., et al. Mol. Vis. 16, 1601-1609 (2010) :
Paliwal, P., et al. Mol. Vis. 16, 1429-1438 (2010) :
Yang, J., et al. Mol. Vis. 16, 1186-1193 (2010) :

TGFBI Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)