

# **TRIM25 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16857c

## **Specification**

#### TRIM25 Antibody (Center) - Product Information

Application WB,E Q14258 Primary Accession NP 005073.2 Other Accession Reactivity Human Host Rabbit Clonality **Polyclonal** Isotype Rabbit Ig Calculated MW 70973 Antigen Region 277-306

TRIM25 Antibody (Center) - Additional Information

#### **Gene ID 7706**

#### **Other Names**

E3 ubiquitin/ISG15 ligase TRIM25, 632n3, Estrogen-responsive finger protein, RING finger protein 147, Tripartite motif-containing protein 25, Ubiquitin/ISG15-conjugating enzyme TRIM25, Zinc finger protein 147, TRIM25, EFP, RNF147, ZNF147

#### Target/Specificity

This TRIM25 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 277-306 amino acids from the Central region of human TRIM25.

#### **Dilution**


WB~~1:1000

### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



TRIM25 Antibody (Center) (Cat. #AP16857c) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the TRIM25 antibody detected the TRIM25 protein (arrow).

# TRIM25 Antibody (Center) - Background

The protein encoded by this gene is a member of the

tripartite motif (TRIM) family. The TRIM motif includes three

zinc-binding domains, a RING, a B-box type 1 and a B-box type 2,

and a coiled-coil region. The protein localizes to the cytoplasm.

The presence of potential DNA-binding and dimerization-transactivation domains suggests that this protein may

act as a transcription factor, similar to several other members of

the TRIM family. Expression of the gene is upregulated in response

to estrogen, and it is thought to mediate estrogen actions in

breast cancer as a primary response gene.

## TRIM25 Antibody (Center) - References





cycles.

### **Precautions**

TRIM25 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

TRIM25 Antibody (Center) - Protein Information

Name TRIM25

Synonyms EFP

{ECO:0000303|PubMed:8248217}, RNF147

#### **Function**

Functions as a ubiquitin E3 ligase and as an ISG15 E3 ligase (PubMed:<a href="http://www.uniprot.org/citations/16352599" target="\_blank">16352599</a>). Involved in innate immune defense against viruses by mediating ubiquitination of DDX58 and IFIH1 (PubMed:<a href="http://www.uniprot.org/citations/17392790" target="\_blank">17392790</a>, PubMed:<a href="http://www.uniprot.org/citations/17392790</a>

PubMed:<a href="http://www.uniprot.org/ci tations/30193849"

target="\_blank">30193849</a>). Mediates 'Lys-63'-linked polyubiquitination of the DDX58 N-terminal CARD-like region and may play a role in signal transduction that leads to the production of interferons in response to viral infection (PubMed:<a href="http://www.uniprot.org/citations/1739279">href="http://www.uniprot.org/citations/1739279</a>)" target="\_blank">17392790</a>, PubMed:<a href="http://www.uniprot.org/citations/23950712"

target="\_blank">23950712</a>). Mediates 'Lys-63'- linked polyubiquitination of IFIH1 (PubMed:<a href="http://www.uniprot.org/c itations/30193849"

target="\_blank">30193849</a>). Promotes ISGylation of 14-3-3 sigma (SFN), an adapter protein implicated in the regulation of a large spectrum signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/16352599"


target=" blank">16352599</a>,

PubMed: <a href="http://www.uniprot.org/ci tations/17069755"

target="\_blank">17069755</a>). Mediates estrogen action in various target organs (PubMed:<a href="http://www.uniprot.org/c itations/22452784"

target="\_blank">22452784</a>). Mediates the ubiquitination and subsequent proteasomal degradation of ZFHX3

Dai, H., et al. Oncol. Rep. 23(3):795-799(2010) Zhao, J., et al. BMC Med. Genet. 11, 96 (2010): Gack, M.U., et al. Cell Host Microbe 5(5):439-449(2009) Ludwig, S., et al. Cell Host Microbe 5(5):420-421(2009) Dai, H., et al. Oncol. Rep. 21(2):395-401(2009)





(PubMed:<a href="http://www.uniprot.org/c itations/22452784" target="\_blank">22452784</a>). Plays a role in promoting the restart of stalled replication forks via interaction with the KHDC3L-OOEP scaffold and subsequent ubiquitination of BLM, resulting in the recruitment and retainment of BLM at DNA replication forks (By similarity).

#### **Cellular Location**

Cytoplasm. Cytoplasm, Stress granule. Nucleus {ECO:0000250|UniProtKB:Q61510}

#### **Tissue Location**

Expressed in breast tumors (at protein level). Ubiquitous.

# **TRIM25 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture