

CARM1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16980c

Specification

CARM1 Antibody (Center) - Product Information

Application	WB, E
Primary Accession	Q86X55
Other Accession	Q4AE70 , Q9WVG6 , NP_954592.1
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	347-375

CARM1 Antibody (Center) - Additional Information

Gene ID 10498

Other Names

Histone-arginine methyltransferase CARM1, 211-, Coactivator-associated arginine methyltransferase 1, Protein arginine N-methyltransferase 4, CARM1, PRMT4

Target/Specificity

This CARM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 347-375 amino acids from the Central region of human CARM1.

Dilution


WB~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



CARM1 Antibody (Center) (Cat. #AP16980c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the CARM1 antibody detected the CARM1 protein (arrow).

CARM1 Antibody (Center) - Background

Protein arginine N-methyltransferases, such as CARM1, catalyze the transfer of a methyl group from S-adenosyl-L-methionine to the side chain nitrogens of arginine residues within proteins to form methylated arginine derivatives and S-adenosyl-L-homocysteine. Protein arginine methylation has been implicated in signal transduction, metabolism of nascent pre-RNA, and transcriptional activation (Frankel et al., 2002 [PubMed 11724789]).

CARM1 Antibody (Center) - References

Gao, X., et al. J. Cell. Biochem. 110(1):162-170(2010)
Carascosa, S., et al. Genes Dev. 24(7):708-719(2010)
Kim, Y.R., et al. BMC Cancer 10, 197 (2010) :
Ito, T., et al. BMC Dev. Biol. 9, 47 (2009) :

Precautions

CARM1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Haiman, C.A., et al. BMC Cancer 9, 43 (2009) :

CARM1 Antibody (Center) - Protein Information

Name CARM1

Synonyms PRMT4

Function

Methylates (mono- and asymmetric dimethylation) the guanidino nitrogens of arginyl residues in several proteins involved in DNA packaging, transcription regulation, pre-mRNA splicing, and mRNA stability. Recruited to promoters upon gene activation together with histone acetyltransferases from EP300/P300 and p160 families, methylates histone H3 at 'Arg-17' (H3R17me), forming mainly asymmetric dimethylarginine (H3R17me2a), leading to activate transcription via chromatin remodeling. During nuclear hormone receptor activation and TCF7L2/TCF4 activation, acts synergically with EP300/P300 and either one of the p160 histone acetyltransferases NCOA1/SRC1, NCOA2/GRIP1 and NCOA3/ACTR or CTNNB1/beta-catenin to activate transcription. During myogenic transcriptional activation, acts together with NCOA3/ACTR as a coactivator for MEF2C. During monocyte inflammatory stimulation, acts together with EP300/P300 as a coactivator for NF-kappa-B. Acts as coactivator for PPAR, promotes adipocyte differentiation and the accumulation of brown fat tissue. Plays a role in the regulation of pre-mRNA alternative splicing by methylation of splicing factors. Also seems to be involved in p53/TP53 transcriptional activation. Methylates EP300/P300, both at 'Arg-2142', which may loosen its interaction with NCOA2/GRIP1, and at 'Arg-580' and 'Arg-604' in the KIX domain, which impairs its interaction with CREB and inhibits CREB-dependent transcriptional activation. Also methylates arginine residues in RNA- binding proteins PABPC1, ELAVL1 and ELAV4, which may affect their mRNA- stabilizing properties and the half-life of their target mRNAs.

Cellular Location

Nucleus. Cytoplasm. Note=Mainly nuclear during the G1, S and G2 phases of the cell cycle. Cytoplasmic during mitosis, after breakup of the nuclear membrane

Tissue Location

Overexpressed in prostate adenocarcinomas and high- grade prostatic intraepithelial neoplasia

CARM1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)