

**CTNNBL1 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP17056a**

### Specification

**CTNNBL1 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8WYA6</a>
Other Accession	<a href="#">NP_110517.2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	65173
Antigen Region	16-43

**CTNNBL1 Antibody (N-term) - Additional Information**

**Gene ID** 56259

**Other Names**

Beta-catenin-like protein 1,  
Nuclear-associated protein, NAP, Testis  
development protein NYD-SP19, CTNNBL1,  
C20orf33

**Target/Specificity**

This CTNNBL1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 16-43 amino acids from the N-terminal region of human CTNNBL1.

**Dilution**


WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



CTNNBL1 Antibody (N-term) (Cat. #AP17056a) western blot analysis in WiDr cell line lysates (35ug/lane). This demonstrates the CTNNBL1 antibody detected the CTNNBL1 protein (arrow).

**CTNNBL1 Antibody (N-term) - Background**

The protein encoded by this gene contains an acidic domain, a putative bipartite nuclear localization signal, a nuclear export signal, a leucine-isoleucine zipper, and phosphorylation motifs. In addition, the encoded protein contains Armadillo/beta-catenin-like repeats, which have been implicated in protein-protein interactions. Although the function of this protein has not been determined, the C-terminal portion of the protein has been shown to possess apoptosis-inducing activity. [provided by RefSeq].

**CTNNBL1 Antibody (N-term) - References**

Ichikawa, S., et al. J. Bone Miner. Res. 25(8):1821-1829(2010)  
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253

**Precautions**

CTNNBL1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

(2010) :

Badiglian Filho, L., et al. *Oncol. Rep.* 21(2):313-320(2009)

Andreasen, C.H., et al. *BMC Med. Genet.* 10, 17 (2009) :

Vogel, C.I., et al. *BMC Med. Genet.* 10, 14 (2009) :

**CTNNBL1 Antibody (N-term) - Protein Information**

**Name** CTNNBL1

**Synonyms** C20orf33

**Function**

Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing. Participates in AID/AICDA-mediated Ig class switching recombination (CSR). May induce apoptosis.

**Cellular Location**

[Isoform 1]: Nucleus.

**Tissue Location**

Widely expressed with highest levels in skeletal muscle, placenta, heart, spleen, testis and thyroid

**CTNNBL1 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)